



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Colloid Transport in Porous Media: A Review of Classical Mechanisms and Emerging Topics

Citation for published version:

Molnar, IL, Pensini, E, Asad, MA, Mitchell, CA, Nitsche, LC, Pyrak-nolte, LJ, Miño, GL & Krol, MM 2019, 'Colloid Transport in Porous Media: A Review of Classical Mechanisms and Emerging Topics', *Transport in Porous Media*, pp. 1-28. <https://doi.org/10.1007/s11242-019-01270-6>

Digital Object Identifier (DOI):

[10.1007/s11242-019-01270-6](https://doi.org/10.1007/s11242-019-01270-6)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Transport in Porous Media

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Colloid Transport in Porous Media: A Review of Classical Mechanisms and Emerging Topics

Ian L. Molnar^{1,*}, Erica Pensini², Md Abdullah Asad¹, Chven A. Mitchell³, Ludwig C. Nitsche⁴,
Laura J. Pyrak-Nolte³, Gastón L. Miño^{5,6}, Magdalena M. Krol¹

¹Lassonde School of Engineering, York University, Toronto, ON M3J 1P3, Canada

²School of Engineering, University of Guelph, Guelph, ON, Canada N1G 2W1

³College of Science, Purdue University, West Lafayette, IN, 47906, USA

⁴College of Engineering, University of Illinois, Chicago, IL, 87185 USA

⁵Inst. de I+D en Bioingeniería y Bioinformática (IBB), CONICET-UNER, Oro Verde, ER, Argentina

⁶LAMAE, Facultad de Ingeniería, Univ. Nac. de Entre Ríos, Oro Verde, Argentina

*Corresponding author at: Department of Civil Engineering, York University, Toronto, ON M3J 1P3, Canada. Email address: imolnar@yorku.ca. ORCID: <https://orcid.org/0000-0002-6610-4975>

Acknowledgments

The work related to the experiments on swarm transport in fractured and porous media, simulation runs, and analysis, was supported by the U. S. Department of Energy, Office of Science, Office of Basic Energy Sciences, the Geosciences Research Program, under Award Number (DE-FG02-09ER16022). We also acknowledge the support of the Natural Sciences and Engineering Research Council of Canada (NSERC). The authors would also like to acknowledge Dr. Sarah A. Charron for her helpful insights into the role of colloids in medical sciences.

Abstract:

To celebrate the tenth anniversary of Interpore, we present an interdisciplinary review of colloid transport through porous media. This review aims to explore both classical colloid transport and topics that fall outside that purview and thus offer transformative insights into the physics governing transport behaviour. First, we discuss the unique colloid characteristics relative to molecules and larger particles. Then, the classical advection-dispersion-filtration models (both conceptual and mathematical) of colloid transport are introduced as well as anomalous transport behaviours. Next, the forces of interaction between colloids and porous

media surfaces are discussed. Fourth, applications that are interested in maximizing the transport of colloids through porous media is considered. Then the concept of motile, active, biocolloids are introduced and, finally, colloid swarming as a newly-recognized mode of transport is summarized.

1. Introduction

Colloids are nano- or micrometer-sized particles (traditionally defined as between 1nm and 10 μm [1]) which include engineered nanomaterials, viruses, bacteria, fine powders, proteins, and finely textured soil particles (e.g., clay). The transport of colloids through porous media is relevant to a wide range of fields such as medical sciences, industrial manufacturing, geosciences and environmental engineering.

Many industrial processes employ porous media filters to either recover valuable colloidal materials or to purify liquid streams of unwanted materials [1, 2]. Vehicles rely on porous filters to ensure that operation of the fuel, transmission and brake systems are not compromised by colloids [3]. The recovery of valuable metals from waste products, such as catalytic converters found on vehicles or discarded batteries, utilize filtration with porous media as a key step in the recovery process [4-6]. Air filtration in both industrial and consumer products uses porous media to remove airborne colloids [7, 8]. In addition, many systems use emulsions which are an important class of industrial colloids [9].

In medical sciences, colloids such as albumin migrate through the body via porous blood vessels and play a vital role in maintaining human health. While the porous circulatory system is permeable to many molecules, proteins are too large to diffuse across the blood-tissue barrier [10]; this gives albumin the ability to maintain oncotic pressure, i.e. the osmotic pressure that prevents serum from leaking out of capillaries and causing a condition known as edema [11, 12]. In addition, innovative cancer treatment technologies exploit colloid transport behaviour to specifically target cancerous cells [13].

In the geosciences and environmental engineering, there is significant interest in predicting how far colloids such as pathogenic bacteria and viruses travel through soils or other porous media [14-18]. Improper design or maintenance of water treatment filtration systems can result in pathogens entering the water distribution system and cause widespread illness [19]. In

addition, communities utilizing groundwater as a drinking water source rely on the natural soil to act as a filter. There are numerous instances that illustrate how disease outbreaks can occur in communities when pathogens are not adequately filtered during transport through soil [20-24]. Natural geological colloids (e.g., clay particles) are also capable of enhancing the transport of dissolved compounds through porous media as relatively immobile contaminants in the subsurface environment may sorb onto the colloid's surface and be transported further-than-expected distances [25-30].

A large volume of excellent research exists on the transport of colloids through porous media. Field-specific review papers synthesize the current knowledge regarding colloid fate and transport in porous media for medical sciences [10, 31-33], geosciences and environmental engineering [14, 16, 18, 34-40], and industrial manufacturing [2, 3, 41-43]. Other reviews have focused on specific colloid types such as metallic nanoparticles [44-46], or on specific mechanisms such as surface forces [47-51].

The discipline and mechanism-specific reviews currently available in the literature have proven invaluable in synthesizing research and outlining future research needs within their respective areas. However, colloid science researchers would benefit from an interdisciplinary resource that ties together the transport mechanisms that govern colloid transport across a range of scales and disciplines. This review paper aims to fill this gap by providing an introductory resource on the fate and transport of colloids in porous media.

2. Unique Characteristics of Colloids

Colloids have traditionally been defined on the basis of their size. However, differing size ranges appear throughout the literature, depending on the properties of the systems examined. There is no strict definition on what separates a colloid from a molecule (many macro-molecules are considered colloids), or from larger particles. Historically, this delineation has been the subject of some controversy, especially within the biological sciences [e.g., 52]. As a result, it is more useful to define colloids by the properties which influence transport behaviour. Compared to larger particles, colloids exhibit higher rates of diffusive flux (e.g., Brownian motion), higher reactivity due to higher surface areas and are more strongly influenced by short range surface forces. Conversely, colloids possess more deterministic trajectories than smaller molecules, and

their forces of interaction (while based on the same fundamental forces that exist between atom-atom and molecule-molecule pairings) can be larger and decay more slowly across distances than smaller molecules [53]. As diffusivity is inversely proportional to fluid viscosity [54], defining colloids on the basis of transport properties suggests that larger aerosols may possess the same diffusivity of smaller aerosols, but not the same specific surface area or reactivity.

2.1. Brownian Motion

Colloidal diffusion manifests as the movement of colloids from regions of high particle concentrations to regions of low particle concentrations, similar to the molecular diffusion of individual molecules. However, unlike individual molecules, colloids diffuse due to their random walk behaviour, known as ‘Brownian motion’, which arises due to the large number of collisions that occur between colloids and molecules in the surrounding fluid. The magnitude of Brownian motion increases with decreasing particle size [40].

The Brownian motion of colloids has implications for colloid transport, even in scenarios where the overall diffusive flux is minor relative to the rate of advective transport. Brownian motion alters the trajectories of individual colloids during transport and, ultimately, impacts how far they may be transported through porous media. The trajectory of a single colloid in a flowing fluid can be described using a combination of deterministic (external and hydrodynamic/aerodynamic forces) and stochastic (Brownian force) forces [55, 56].

During transport through a porous medium, a colloid’s partially deterministic trajectory causes it to follow flow streamlines through the medium. The Brownian motion aspect of the colloid’s trajectory allows it to cross flow streamlines and contact surfaces, or other colloids, that were not on the colloid’s initial trajectory [40] (illustrated in Figure S1 of the supplementary information).

2.2. Colloid Surface Area

The specific surface area of particles (i.e. the surface area of a substance per unit mass) increases with decreasing particle diameter. For a single spherical colloid, its surface area (A_s) is given as $A_s = \pi d_p^2$, while its mass (M_s) is $M_s = \frac{1}{6} \pi d_p^3 \rho_s$, where d_p is the colloid’s diameter and

ρ_s is the colloid's density. The specific surface area (SSA) of a spherical colloid can be described as:

	$SSA = \frac{6}{d_p \rho_s}$	(2.1)
--	------------------------------	-------

This indicates that high SSA is another unique colloid characteristic and may result in colloids being highly reactive. Chemical reactions between colloids and solutes occur on the surface of the colloid [57-59], thus as particle size decreases the total area available for reaction increases, increasing its reactivity [60] and capacity for surface absorption [61]. Reactions may alter the small-scale forces of interaction that govern colloid retention and agglomeration (defined as an assemblage of colloids, not to be confused with pre-nucleation aggregates [62]).

3. Advective-Dispersive Transport and Filtration Behaviour

Colloid transport through porous media is classically described at the continuum-scale (i.e., macroscopic-scale) using the classic advection-dispersion partial differential equation (ADE). The ADE typically takes the form described by Eq. 3.1:

	$\frac{\partial C}{\partial t} \phi = -\nabla \cdot (\vec{v}C)\phi + \nabla \cdot (\overline{D_{disp}} \nabla C)\phi - \rho_b \frac{\partial S}{\partial t}$	(3.1)
--	--	-------

where C is the concentration of colloids suspended in fluid ($M \cdot L^{-3}$), t is time (t), ϕ is the medium's porosity, \vec{v} is the pore velocity vector ($L \cdot t^{-1}$), $\overline{D_{disp}}$ is dispersion tensor ($L^2 \cdot t^{-1}$), ρ_b is the bulk density of the medium ($M \cdot L^{-3}$) and S is the mass of colloids retained (i.e. filtered) per mass of porous medium ($M \text{ colloids} \cdot M^{-1} \text{ soil}$). Size exclusion (i.e. colloids unable to enter smaller pores by virtue of their size) may change the effective porosity of the medium and result in colloids travelling at faster-than-expected velocities [63]. The dispersion coefficient represents a non-physical fitting parameter that changes with scale [64]; while this indicates that Eq. 3.1 is not a completely accurate representation of colloid transport, it remains a useful simplification that is utilized for colloid transport through geologic media [40], biological tissues [65], and fibrous filters [3], among others.

Generally, colloids are transported in response to gradients of thermodynamic quantities (e.g., colloid diffusion occurs in response to a colloid concentration gradient). As a result, there are numerous other modes of colloid transport from different thermodynamic quantities not

shown in Eq. 3.1 such as: Diffusiophoresis and chemotaxis in which colloids move in response to a solute's gradient [66, 67], electrophoresis [68], magnetophoresis [69], and thermophoresis [70]. In addition, self-propelled colloids are the subject of research for several fields, while many utilize the gradients discussed above [71, 72], motile biocolloids can be mechanically driven via appendages such as flagella [73, 74].

The small-scale interactions between colloids and porous media surfaces govern how far colloids travel via a process known as filtration. The standard conceptual model of filtration is that a porous medium is comprised of pore spaces (i.e., void/fluid phase) and 'collectors' which are discrete units of the solid phase [75]. Colloids are transported through a medium's pore space via advection and dispersion/diffusion until they enter a collector's near-surface zone (typically on the order of nanometers from a collector's surface). Colloids are brought into this near surface zone either when the streamline brings the colloid close to the surface (interception), gravity forces causes the colloid to approach a collector, when Brownian motion carries the colloid across streamlines, or when inertial forces carry the impinging colloid across streamlines and toward the collector [56, 76-80].

The small-scale forces of interaction in the near-surface zone (discussed in section 4) impact whether the colloids are 'retained' by the collector or continue travelling. This process in which colloids are removed from the fluid phase is known as deep bed filtration, which is classically described as a first-order kinetic process with the retention parameter from Eq. 3.1, $\rho_b \frac{\partial S}{\partial t}$, rewritten in terms of:

	$\rho_b \frac{\partial S}{\partial t} = k_{ret} C \phi$	(3.2)
--	---	-------

where k_{ret} is the first-order kinetic retention parameter (t^{-1}). Depending on the colloid-colloid forces of interaction, retained colloids may provide further sites for colloid retention, yielding a filter ripening/clogging effect which can decrease medium permeability (i.e., increases a filter's pressure drop) and increase the rate at which colloids are retained [76, 81-84]. For the special case of colloid transport through biological tissues, filtration is a two-step process: 1) adsorption/desorption on the cell surface, and 2) internalization/uptake into the cell body, both of which are describable via first-order processes [85, 86]. Colloid filtration by biological tissue is

further limited by the number of available binding /reaction sites on an individual cell [87],
conceptually similar to site blocking proposed for geological media [88].

3.1. Colloid Filtration Theory

There is significant interest in developing mechanistic approaches to estimating k_{ret} (Eq. 3.2) to allow a-priori predictions of colloid transport through a porous medium, without relying on empirically determined parameters. The most common approach to mechanistically predicting k_{ret} is colloid filtration theory (CFT), which accounts for the properties of the porous medium, the colloids, the flow regime, and the chemical conditions [56, 75, 76, 80, 82, 89-95]. There are two primary components to modern CFT formulations: 1) a mechanistic model that solves a force/torque balance to estimate the trajectory of a colloid near a collector, and 2) a correlation equation that summarizes the results of the mechanistic model. These results are then upscaled to k_{ret} [40, 96-98]. For a granular medium idealized as a packed bed of Happel Sphere in Cells with diameter d_c , k_{ret} is estimated via Eq. 3.3 [90] where η_o is derived from CFT and represents the fraction of colloids approaching a collector that are retained:

	$k_{ret} = \frac{3}{2} \frac{(1 - \phi)}{d_c} \eta_o v$	(3.3)
--	---	-------

While numerous CFT models exist for granular media in water [56, 80, 90, 91, 99-102], similar models exist for aerosol filtration by fibrous filters which consider flow across a series of cylinder [76, 82, 83, 94].

The accuracy of CFT's prediction of k_{ret} , is governed by the small-scale surface forces that dominate colloid-collector interactions in the near surface zone. In scenarios where the colloid-collector surface forces are attractive (i.e., so-called favorable attachment conditions), each colloid-collector contact event results in retention and ADE/CFT generally provides an excellent match to experimental colloid transport data [56]. However, there are scenarios where colloid-collector surface interactions are repulsive and attachment conditions are unfavorable. Unfavourable attachment conditions are predominant in environmental scenarios, as many types of both geologic media and natural colloids carry negative electrostatic charges [40] which yield repulsive forces of interaction. Under unfavourable attachment conditions CFT models are not accurate. The CFT models predict no colloid retention, as the repulsive surface forces prevent

colloid retention [103]. Instead, experimental observations of colloid retention in unfavorable conditions are widespread throughout colloid transport literature [27, 35, 104, 105] which has led to the development of an additional CFT parameter, α , termed attachment efficiency. The attachment efficiency parameter is defined as the fraction of colloids that contact the collector and are retained. Attachment efficiency is incorporated into descriptions of CFT's k_{ret} in Eq. 3.3 as follows [103, 104, 106]:

	$\eta_o = \eta\alpha$	(3.4)
--	-----------------------	-------

where η is the fraction of colloids approaching the collector that contact the collector (i.e., enter its near-surface zone). Understanding the factors that impact α , and developing predictive models for it, has been the subject of a large number of studies [35, 106-112]. However, no method has successfully demonstrated an ability to predict α a-priori, or fully describe the impacting factors and, as a result, generally remains a fitting parameter.

3.2. Anomalous Transport Behaviour

The first-order retention rate in Eq. 3.2 predicts an exponential retention profile of colloids (i.e., the concentration of retained colloids decreases exponentially with distance). In addition, the ADE predicts a breakthrough curve with a flat plateau and symmetrically sharp increases and decreases in colloid concentration during breakthrough and elution.

Observations of anomalous colloid transport behaviour are widespread. Experimental retention profiles in unfavorable conditions are reported to be hyper-exponential, linearly decreasing or non-monotonic [113-121]. Breakthrough curves may be non-symmetric, a slow increase in concentration over time [122-124], a slow decrease (i.e., filter ripening\clogging) [125], or there may be a slow decrease in colloid concentration during elution (i.e., tailing) (illustrated in Figure S2 in the supplementary information) [126-130]. The application of α to calculate k_{ret} in unfavorable conditions does not resolve the anomaly between experimentally observed anomalous behaviours and the behaviour predicted by the ADE/CFT approach. The predicted retention and breakthrough behaviours are due to the use of k_{ret} as a first-order irreversible retention rate coefficient, applying α as a scaling coefficient for k_{ret} will not change its reaction order or predicted behaviours.

Several theories have been proposed to explain these anomalous behaviours. Non-exponential retention profiles have been explained with theories such as fast and slow colloid retention rates [118], and dual porosities such as mobile and immobile pore space in geologic media [131, 132] or vascular and interstitial pore spaces in biological media [133]. Transport in unfavorable conditions have been hypothesized to be a result of retention at small distances from the collector surface in an energetically favorable region known as the secondary energy minimum [105, 116, 134] or by incorporating zones of discrete micro- and nano-sized zones of favorable attachment conditions on the collector surface [135, 136] and heterogeneous binding sites on cell membranes [137]. In addition, reversible retention and colloid re-entrainment into the bulk pore fluid is considered necessary to produce the observed tailing behaviours [138-140]. However, the macroscopic rate coefficients employed by these models are empirical and can only be determined based on experimental data.

Liquid aerosols also exhibit anomalous transport behaviour. Numerous studies have reported that clogging increases the penetration of liquid aerosols through the filter [141-143], contrary to the decreased penetration observed from solid aerosols. This has been attributed to the aerosol liquid wetting filter fibers [141, 142] as different behaviour is observed under non-wetting conditions [144].

Each CFT model is only valid for a specific porous media and a narrow range of physicochemical conditions [40]. The specific transport mechanisms (i.e., formulation of Eq. 3.1) should be chosen based on desired outcomes and properties of the system; systems with multiple porosities (e.g., biological media) would benefit from a multi-porosity description [133], and weakly retained colloids would require desorption/re-entrainment mechanisms [85], or incorporating transport via gradients of different thermodynamic quantities [71]. In the cases of anomalous transport behaviour, purely mathematical models such as continuous time random walk (CTRW) can accurately match experimental observations [145], but yield little insight into the governing mechanisms. Given the wide variety of models, their varying ranges of applicability and their successes in describing both classic and anomalous colloid transport, the choice of conceptual or computational model to assess colloid transport should not be arbitrary, especially with regards to colloid-collector forces of interaction.

4. Forces of Interaction

As discussed in section 3, the forces of interaction determine how colloids are filtered out of solution and retained by the porous media. As a result, the forces that govern colloid-colloid and colloid-porous media interactions are key to understanding how colloids are transported through porous media and to implementing mechanistic models such as colloid filtration theory.

4.1. Forces of interaction before contact: the DLVO theory

The forces of interaction between colloid-porous media and colloid-colloid interactions are traditionally described using the Derjaguin-Landau-Verwey-Overbeek (DLVO) theory [146].

As colloid-porous media and colloid-colloid interactions are governed by the same forces of interaction, the terms bodies and particles are employed interchangeably throughout this discussion to generically refer to both colloids and porous media. In addition, the term surfaces is used throughout to refer to either colloidal or porous media *surfaces*.

The DLVO model theorizes that the electric charge on surfaces creates an electric potential difference between the surface and the solution, hence attracting oppositely charged ions in solution (counterions). The counterions are simultaneously attracted to the surface (e.g. by electrostatic forces) and dispersed (e.g. thermal Brownian motion). At equilibrium, a counterion concentration gradient exists around the charged surface, with the counterions closest to the surface sensing the strongest attraction to it. The counterion layer around the surface (electrical double layer) can be divided into two layers: 1) the compact Stern layer, mainly subject to the attraction between the counter-ion and the surface, and 2) the shear plane (i.e. the outermost surface), mainly subject to the effect of Brownian motion.

The zeta-potential (i.e. the electric potential at the shear plane) determines the electrostatic interactions between a colloid and other bodies (e.g. porous media). According to the DLVO theory, the zeta-potential (ζ) ($\text{M}\cdot\text{L}^2\cdot\text{A}^{-1}\cdot\text{t}^{-3}$; A denotes current) is related to the charge Q ($\text{A}\cdot\text{t}$) of the shear plane as follows (Eq. 4.1):

	$\zeta = \frac{4\pi\delta Q}{\varepsilon}$	(4.1)
--	--	-------

where δ is the thickness of the diffusive layer (L) and ε is the dielectric constant of the liquid (water).

The double layer (DL) theory has been widely employed to describe the forces of interactions between bodies in fluid. DL forces ($M \cdot L \cdot t^{-2}$) between a sphere and a plane may be for instance expressed as described in Eq. 4.2 [147]:

	$F_{electro} = 128\pi r \gamma_{sphere} \gamma_{plane} n k_b T \kappa^{-1} \exp(-\kappa d)$	(4.2)
--	---	-------

where: r = radius of the sphere (L), γ_{sphere} and γ_{plane} are lumped parameters, $\gamma_i = \tanh\left(\frac{e\psi_i}{4k_b T}\right)$ where ψ_i is the surface potential (V) and e is the charge of an electron. n = concentration of ions in the bulk solution ($M \cdot L^{-3}$), κ^{-1} = Debye's length (L) and d = distance between the surface of the sphere and the plane (L). The DL force can be attractive or repulsive, depending on the charge of the interacting bodies (e.g. colloids and porous media).

The Debye length measures the range of the double layer. The DLVO theory fails in cell biology applications, when the Debye length is particularly small ($< 8 \text{ \AA}$; where \AA denotes Angstrom), in micellar chemistry applications in which the salt concentrations are high, or in ionic microemulsions or polyelectrolytes systems, for which the screening length is comparable to the hydrated ion sizes [148].

In addition to the DL force, the DLVO theory accounts for van der Waals (VdW) forces and, later, Born forces. VdW forces are always attractive in aqueous media. They represent multipole- multipole interactions and can be classified as follows [149, 150]: 1) Keesom interactions, i.e. the interactions between charges (in the case of molecular ions) and between permanent multipoles; 2) Debye interactions, i.e. the interactions between a permanent multipole with an induced multipole and; 3) London interactions (or dispersion forces), which act between any pair of molecules, regardless of their symmetry.

The VdW interactions are a function of d^{-n} and become infinite when d is zero, whereas DL forces remains finite or rises more slowly than VdW forces [150]. A complete derivation of the expression for VdW forces can be found elsewhere [150]. The VdW forces ($M \cdot L \cdot t^{-2}$) between a sphere and a plane are described by [151]. At short separation distances, the magnitude of VdW forces is always greater than that of DL forces, as described earlier. Therefore, when only VdW and DL forces are considered, the DLVO model predicts infinite attraction at zero separation distance (termed 'primary energy minimum'). Therefore, the DLVO model was expanded to include Born repulsive forces [152] and has been expressed by [151].

The DLVO theory was developed to describe the interactions between hard surfaces, and it is hence not well-suited to describe the interactions between soft charged colloids (i.e., colloids with an ion-permeable surface layer [153]). A new theory was developed by Ohshima to describe the interactions between bodies with a hard core and a soft outer shell [153]. However, the Ohshima model does not account for steric and hydration forces, the importance of which is outlined in section 4.2.

4.2. Non-DLVO forces of interaction before contact: steric, bridging and hydration forces

Steric forces stem from the presence of chains dangling from a surface, e.g. polymer chains attached to colloid surfaces. When polymers overlap, they exert a repulsive osmotic force, the magnitude of which depends on the polymer chemistry and density on the surfaces, on the way they are attached to the surfaces (i.e. if they are physically adsorbed or grafted onto the colloid surface) and on the characteristics of the solvent. In the presence of polymers, bridging forces may also arise because polymers protruding from the surface of a colloidal particle can stick to the surface of a second colloidal particle.

Hydration forces are also not accounted for in the DLVO theory. Hydration forces originate from water structuring at solid-liquid interfaces. In the bulk fluid, hydrogen bonds between water molecules dictate the structure of water. Foreign solutes or colloids introduced in solution affect the hydrogen bonding network because they exclude water molecules, acting as cavities [154]. Small solutes can squeeze between water molecules without breaking any hydrogen bonds. Conversely, colloids break the hydrogen bonds and disrupt the structure of water around the surface of the immersed body [154]. The changes induced in the structure of water depend on the characteristics of the solid surface. In the case of hydrophilic bodies, water molecules become tightly bound onto the surface, preventing the approach of other bodies and producing repulsive interactions [150].

4.3. Relevance of the forces of interaction on colloid transport in geological media

The effect of the forces of interaction on colloid transport in geological media is three-fold: 1) they affect the aggregation of colloids; 2) they determine the rheological behavior of fluids carrying the colloids; 3) they determine if colloids are retained within geological media or migrate in the subsurface. The aggregation behavior of colloidal particles (and particularly of nano-colloidal particles) strongly affects subsurface transport [155, 156]. Attractive forces induce agglomeration, which inhibits transport by clogging the pores, whereas repulsive forces between colloids ensure adequate dispersion and promote their transport. The interplay between hydrodynamic forces and the forces of interaction between colloidal particles and geological media (soil grains or fractured rocks) is also paramount in determining colloid transport [157-164]. Non-DLVO forces play a particularly relevant role when the interacting surfaces are strongly hydrated such as silica and iron [158, 159, 161], when they are soft [159-161, 165], or in the presence of extracellular polymeric substances produced by bacteria [166].

4.4. Forces of interaction and colloid transport in living organisms

Similar to colloid transport in geological media, colloid transport in the mammal and frog body is influenced by the electric charge of the colloids. The importance of electrostatic interactions has been recognized for positive charged colloids interacting with the walls of the gastrointestinal tract [167] and in the context of drug delivery in tumor tissues [168].

Attractive van der Waals forces, steric and depletion forces (e.g. hydrophobic forces), hydration forces (in the case of hydrophilic colloids), and polymer bridging also play a role in the interactions between nanocolloids and circular macromolecules and biological surfaces [169, 170]. The analysis of the interactions between colloids and biological surfaces in the body is complex because the cells are deformable, their surface is heterogeneous, and they are active [169]. Cells can transport ions and secrete molecules, hence modifying the surface of the colloids they interact with [169]. Nanocolloids in contact with biological fluids become coated with proteins, which can change conformation and lead to either increased affinity between proteins and surfaces or to the exposure of new epitopes (i.e. the part of an antigen molecule to which an antibody attaches itself) [169]. Moreover, specific receptor–ligand interactions can exist and

cells can incorporate bound colloids (endocytosis), further increasing the complexity of colloidal transport in the body [169].

5. Colloid Injection into Porous Media

Many applications of colloid transport in porous media are focused on filtering colloids out of fluids (e.g., potable water filtration), however there is a unique class of applications that are interested in deliberately injecting colloids into porous media. The fate and transport of these injected colloids governs their efficacy, as result there is significant interest in manipulating colloid behaviour in their respective porous environment to maximize transport. These applications include drug delivery into biological tissues [171], enhancing contrast in specific tissues for medical imaging [172], injecting reactive colloids (e.g., nano-zero valent iron; nZVI) [44, 58, 173] or biocolloids [174, 175] into geologic media to remediate soil and groundwater resources, utilizing colloid-sensors for subsurface sensing [176-178] and generating emulsions in-situ for oil recovery [179]. These disparate applications are governed by a similar set of mechanisms (discussed in section's 3 and 4) and significant opportunity exists for cross-disciplinary learning.

5.1. Stabilizers and surface coatings

Engineered nanoparticles are notoriously unstable (i.e., subject to agglomeration and sedimentation) which can limit mobility [180], inhibit cell uptake in biological tissues [181-184] and reduce colloid reactivity [185]. Colloid flocculation/agglomeration is typically an irreversible process (i.e., kinetically controlled) [186] when colloids interact in the primary energy well; however, in certain conditions colloids may interact in the secondary energy minimum, and agglomeration instead becomes a reversible process [187]. Stabilizers and surface coatings prevent attachment in the primary minimum with repulsive electro-steric forces, in these scenarios reversible attachment governs agglomeration behaviour [185]. As a result, there is a large body of research exploring methods to minimize aggregation and maximize mobility, reactivity and cell uptake in porous media [e.g., 185, 188].

Different kinds of stabilizers and surface coatings have been employed to increase the performance (i.e. stability and reactivity) of nZVI [189, 190] and biomedical sensing and nano-

drug delivery [e.g., 191, 192]. For biocolloid applications there is a strong focus on utilizing surfactants to reduce retention by the porous medium [193, 194]. Many of the surfactants studied for reducing bacterial retention [e.g., 195, 196] are similar to those employed in medical and nZVI stabilization, indicating significant potential for cross-disciplinary collaboration.

5.2. Injected Colloid Mobility

Many colloid injection applications report moderate mobility, with transport distances of nZVI and biocolloids ranging from less than 1m to 6m [193, 197-200] and limited penetration of nano-drugs into tumor tissues [201, 202]. Manipulating injection velocities is a promising approach to improving mobility. Retention rates have been observed to decrease with increasing velocity in unfavorable deposition scenarios [203, 204] due to increased hydrodynamic drag [205]. As a result, constant flux methods yield greater travel distances compared to constant head injection [173] due to higher injection velocities. Variations in flow velocity are capable of remobilizing retained colloids [206] and potentially improving colloid mobility, yet aside from a pressure pulse technique field trial [207] there is little data available on its efficacy for nZVI injection. This behaviour is not universal to all colloid types, as there is evidence that injecting motile bacteria at high velocities may increase retention [208]. Manipulating velocities is also likely of limited usefulness in scenarios with predominantly favorable attachment conditions, as high flow velocities will be needed to detach colloids from the primary minimum [209]. Manipulating pressure gradients in tumors has had little success in improving colloid transport [e.g., 210].

In addition, mobility can be enhanced using recirculation techniques (e.g., continuously extracting and re-injecting groundwater) or flushing with water or stabilizing solution post-injection [211-213]. Other strategies for improving mobility involve decreasing colloid size via stabilization and synthesis techniques [189], starving biocolloids [214, 215], dynamic size changes [201], or developing biocolloids that either produce biosurfactants or are naturally adhesion resistant [216].

Research focused on chemical locomotion and self-propelled colloids [71, 72, 217] should be considered for aiding colloid mobilization through porous media, either via manipulating gradients, selecting colloids that will self-propel along pre-existing gradients, or designing

colloids with catalytic gradients on their surfaces [e.g., 218]. Additional phenomena such as exclusion zones should be further investigated, as some colloid-surface interactions are capable of generating long range repulsive forces that yield zones up to 100 μm in thickness that are colloid free [219, 220].

6. Particle Swarm Transport in Fractures

Despite decades of colloid transport research, new modes of transport are still being discovered, such as particle swarms which have the potential to transform our understanding of colloid transport through geologic or engineered filters, as well as enable novel approaches to applications that deliberately inject colloids into porous media and seek to maximize their transport. Although the settling speeds of individual colloidal particles is very slow, collective hydrodynamics of colloidal swarms settling under gravity can be much more significant and qualitatively different. Swarms are hypothesized to form in porous media when small, localized clusters of colloids begin to collectively migrate and settle under gravity, such conditions may be found in, among others, slow sand filters and fractured bedrock. In this brief section we explain the origin and scaling of swarm dynamics and indicate how the presence of the surrounding walls of pores or fractures can influence the collective behavior.

A baseline for the behavior of colloidal swarms is provided by the available literature on open-tank conditions [221-228], which is strikingly similar to the behavior of descending vortex rings formed from miscible drops [229-235]. Under conditions of low Reynolds number, colloidal suspensions and miscible drops can be shown to be mathematically equivalent in the gravity-driven Stokes equations [221]: the former represents a kind of approximate discretization of the latter – whether in the laboratory or in a computer simulation. Notwithstanding subtle effects of the granular substructure (fluctuations and modulation of possible influence of fluid inertia [227, 236], a colloidal swarm will therefore behave roughly like a drop of homogeneous miscible liquid with the same density and viscosity as that of the solid-liquid mixture. The analogy of a liquid drop provides the velocity scale: the Stokes settling velocity of an entity of linear dimension d_l and density ρ through an expanse of liquid having density ρ_f and viscosity μ .

	$v_{\text{settling}} = c \frac{(\rho - \rho_f) g d_l^2}{\mu}.$	(6.1)
--	--	-------

The constant c depends only weakly on the shape and viscosity of the entity. Therefore, it is primarily the characteristic diameter-squared dependence that distinguishes the velocities of individual particles versus swarms. For an individual sphere of diameter d_p versus a swarm of diameter d_s and (small) volume fraction ϕ , the respective settling velocities are:

	$v_p = \frac{1}{18} \frac{(\rho_p - \rho_f) g d_p^2}{\mu} \propto d_p^2, \quad v_s = \frac{\phi}{15} \frac{(\rho_p - \rho_f) g d_s^2}{\mu} \propto d_s^2$	(6.2)
--	---	-------

Each particle descends with velocity v_p (Eq. 6.2) relative to its local liquid environment. Since the liquid environment gets dragged downward by the particles settling nearby, a group of particles will settle faster, v_s , than each individual particle. When there are so many particles as to warrant the collective identification as a swarm, then the particles present an additional gravitational force distributed throughout the volume of a grainy pseudo-liquid (i.e., a liquid with an average of the solid- and liquid-phase properties), and the hydrodynamic interactions sum up to produce the characteristic current (a toroidal vortex) inside the swarm, according to the Hadamard-Rybczynski solution [237-239]. Most important, $v_s \gg v_p$ because $(d_s/d_p)^2 \gg \phi^{-1}$ even for dilute suspensions. For a 1 mm diameter swarm composed of 1% concentration by weight of 1 μm spherical colloids (e.g., clay), there would be $\sim 500,000$ particles and $v_s \sim 10,000 v_p$. Thus, colloids in a swarm would be transported with speeds much larger than the settling speed of an individual particle.

The complex shape evolution undergone by colloids swarms and vortex rings is well known [221, 222, 224, 227, 230, 231, 240-243] and includes formation of a ring as an intermediate stage. The most interesting and relevant feature is bifurcation of the swarm into two (or more) smaller swarms. This reduction in size immediately slows down sedimentation according to Eq. 6.2. As the presence of even one planar wall [244] reduces the time/distance to breakup, one would anticipate a similar effect for a fracture which is composed of two walls, i.e. rough surfaces. In addition to fractures, swarms have also been observed in granular porous media. A key component of their transport behaviour through both rough fractures and granular media is the swarm's ability to deform in response to changes in aperture. Figure 1 illustrates the

difference in swarm transport through a rough fracture (Figure 1a) in an impermeable matrix compared to a swarm released in a porous medium composed of spherical grains (Figure 1b,c). For rough fractures, bifurcations occur around asperity peaks (Figure 1a). In the cubic packed porous medium, the swarm expands and contracts as it moves from a pore throat into a pore body and back into a pore throat (Figure 1b). The swarm refocuses at every pore body. The repetitive expansion and contraction produced small oscillations in the velocity-depth curve as the swarm asymptotically approaches a constant velocity. Unlike in the fracture, swarm speed can decrease in a porous media from the repetitive reconfiguration of the swarm.

For fractured porous media (Figure 1c), whether or not an intact swarm is transported across a fracture depends on the volume of the swarm, the aperture of the fracture, and the alignment of pores in the matrices across the fracture. Counter intuitively, large-aperture fractures do not enhance swarm transport between porous matrices. Large-aperture fractures result in a decrease in swarm speed, an increase in the probability of swarm bifurcation, and an increase in the probability of particles remaining in the horizontal fracture (Figure 1c). Large-aperture fractures caused this reduction in speed because a swarm is free to expand laterally in the fracture, increasing the swarm diameter and the drag on the swarm.




Fig. 1 (top row) Time-lapsed composite images and (bottom row) swarm velocity as a function of depth for a swarm in (a) a rough-walled fracture, (b) a cubic packed porous medium, and (c) fractured porous media with the fracture in contact and for a 10 mm aperture, showing swarm deformation in response to changes in confinement geometry.

7. Active Microbiological Colloids

Much of the discussion above has focused on the transport of passive colloids (i.e., colloids incapable of self-propulsion). However, in nature numerous colloids are capable of self-propelled transport (known as active colloids [245-248]) such as sperm inside the female reproductive tract where thousands of cells move through narrow channels in harsh acidic environments [249, 250]. Other examples include the decontamination of waste water through biofilters or slow sand filters. In these systems, biofilms can form at the top of the filter, allowing the retention of both particles and self-propelled microorganisms [251-255]. Active colloids also exist in the field of medicine; the implantation of biocompatible scaffolds allows cells, such as osteoblasts, to colonize porous material and promote tissue regrowth [256]. Even gut microbiota can be considered as a complex system able to contribute to nutrient absorption or immune system responses [257, 258]. These active colloids are not in thermodynamic equilibrium and their study can reveal new mechanisms linking biological processes with physical laws.

7.1. Interactions between swimmers and surfaces or interfaces

Active swimmers may be classified according to the force that they exert on the fluid. Considering far-field hydrodynamics, the simplest model is represented by a dipole force [259-261], and depending on whether the dipole is extensile or contractile, this leads to two categories: pushers and pullers, respectively (illustrated in Figure S3 of the supplementary information). From this point of view, *Escherichia coli* or *Bacillus subtilis* are pushers, and algae such as *Chlamydomonas reinhardtii*, can be considered pullers. These two categories do not fully cover all the possibilities of propulsion at low Reynolds numbers [247, 262-265], but help distinguish two types of effects. Organisms such as *Paramecium* with complex cilia mechanics cannot be described as a dipole [266, 267].

Escherichia coli [268] represents a good model for active colloids[269]. Its body has an ellipsoidal-shape (1 μm diameter and 2 μm length), where the motion is the consequence of the rotation of several helicoidal flagella. Each flagellum is linked to the cell membrane by a nanoscale motor. When all the motors turn counter-clockwise, the flagella rotate in a bundle and push the cell forward. When a motor rotation switches to clockwise, the cell can change direction. When a pusher-like swimmer approaches a solid, it will remain swimming parallel to the wall, producing a long dwelling time or a trapping mechanism at the surface. This phenomenon will generate an increase of bacterial density in vicinities of a solid [270-274]. The flagella or cilia structure can contribute to scattering from the wall [275], despite other evidence that scattering is caused by hydrodynamics [276]. In the absence of flow, interaction with spheres and cylindrical objects is also observed [277, 278]. Puller-type swimmers scatter from curved surfaces, and both hydrodynamic and steric forces are necessary to explain such interactions [279].

7.2. Active Bacteria-Fluid Interactions

It is well known that fluid properties can modify the motility of microorganisms. However, later work has also demonstrated that biologically active particles can modify fluid properties. Living particles in a suspension create a non-equilibrium system where organisms are constantly modifying their surroundings by taking in nutrients and secreting waste [280] such as with fermentation processes, where the production of alcohol reduces fluid density and alters pH. Self-propelled cells can modify fluid rheological properties [281, 282] and exhibit non-Newtonian behavior [283]. Pusher-type swimmers can decrease viscosity [283, 284], flagellar beating as in puller-type swimmers can increase viscosity [285, 286]. Under certain conditions, active colloids can even achieve the characteristic of frictionless superfluids [287, 288].

In addition, flowing fluid can modify bacterial transport behavior [289]. In small channels, high shear forces near walls can re-orient bacteria and result in upstream movement [290, 291]. In addition, active bacteria tend to move from channel centers to the high shear region near lateral solid surfaces [292, 293]. Flow also influences bacterial quorum sensing (i.e. chemical communication) that leads to collective responses and biofilm formation [294].

7.3. Microbiological Transport in Porous Media

In porous media, the flow and shear rate distribution will yield a complex combination of the behaviours discussed above. Bacteria are capable of moving through pores sizes smaller than their body size [295]. Magnetotactic bacteria are able to travel effectively through heterogeneous porous media and overcome relatively high pore water velocities ($\sim 250\mu\text{m/s}$) via magnetotaxis [296]. Bacteria also accumulate on the downstream stagnation points of collectors where flow velocities are smaller than bacteria speed [297]. Despite this work, the influence of motility on retention in porous media remains poorly understood [298]. The complexity of interactions between self-propelled microorganisms and porous media provides substantial opportunities for future research and a valuable avenue for understanding transport processes in porous media and optimizing applications that aim to either limit or maximize colloid mobility in porous media.

8. Conclusions

Despite decades of research into colloid transport through porous media, substantial future research opportunities exist. New mechanisms, phenomena and modes of transport are still continuously being revealed. The origins of many anomalous transport behaviours and colloid-colloid and colloid-collector interactions are still unknown. In addition, new colloid types such as engineered self-propelled colloids and applications such as biomedical and subsurface sensing ensures that colloid transport through porous media will remain a rich and diverse field for many decades yet.

While colloid transport research is conducted across a wide range of disciplines and porous media types, this paper has outlined how the physics governing colloid transport and approaches to solving colloid-related problems, are remarkably similar. For instance, the efficacy of engineered nanoparticles for use in medicine and geosciences are limited by their agglomeration and poor mobility. Similar approaches are being undertaken to improve their delivery, such as manipulating injection conditions and stabilizing particles utilizing electro-steric forces. As a result, cross-discipline collaborations and research will prove invaluable to solving the myriad of outstanding problems in colloid transport through porous media.

References

1. Rhodes, M.J., *Introduction to particle technology*. 2008: John Wiley & Sons.
2. Svarovsky, L., *1 - Introduction to solid-liquid separation*, in *Solid-Liquid Separation (Fourth Edition)*. 2001, Butterworth-Heinemann: Oxford. p. 1-29.

3. Iliev, O., R. Kirsch, Z. Lakdawala, S. Rief, and K. Steiner, *Modeling and Simulation of Filtration Processes*, in *Currents in Industrial Mathematics: From Concepts to Research to Education*, H. Neunzert and D. Prätzel-Wolters, Editors. 2015, Springer Berlin Heidelberg: Berlin, Heidelberg. p. 163-228.
4. Hoffmann, J.E., *Recovery of Platinum-Group Metals from Gabbroic Rocks Metals from Auto Catalysts*. JOM, 1988. **40**(6): p. 40-44.
5. Shin, S.M., N.H. Kim, J.S. Sohn, D.H. Yang, and Y.H. Kim, *Development of a metal recovery process from Li-ion battery wastes*. Hydrometallurgy, 2005. **79**(3): p. 172-181.
6. Rabah, M.A., F.E. Farghaly, and M.A. Abd-El Motaleb, *Recovery of nickel, cobalt and some salts from spent Ni-MH batteries*. Waste Management, 2008. **28**(7): p. 1159-1167.
7. Fisk, W.J., D. Faulkner, J. Palonen, and O. Seppanen, *Performance and costs of particle air filtration technologies*. Indoor Air, 2002. **12**(4): p. 223-234.
8. Biswas, P. and C.-Y. Wu, *Nanoparticles and the Environment*. Journal of the Air & Waste Management Association, 2005. **55**(6): p. 708-746.
9. Bibette, J., F.L. Calderon, and P. Poulin, *Emulsions: basic principles*. Reports on Progress in Physics, 1999. **62**(6): p. 969.
10. Pappenheimer, J.R., *Filtration, Diffusion and Molecular Sieving Through Peripheral Capillary Membranes A Contribution to the Pore Theory of Capillary Permeability*. The American journal of physiology, 1951. **167**(1): p. 13-46.
11. Weil, M.h., R.J. Henning, and V.K. Puri, *Colloid oncotic pressure: clinical significance*. Critical Care Medicine, 1979. **7**(3): p. 113-116.
12. Rackow, E.C., I.A. Fein, and J. Leppo, *Colloid Osmotic Pressure as a Prognostic Indicator of Pulmonary Edema and Mortality in the Critically Ill*. Chest, 1977. **72**(6): p. 709-713.
13. Torchilin, V., *Tumor delivery of macromolecular drugs based on the EPR effect*. Advanced Drug Delivery Reviews, 2011. **63**(3): p. 131-135.
14. Bitton, G. and R.W. Harvey, *Transport of pathogens through soil*, in *Environmental Microbiology*, R. Mitchell, Editor. 1992, Wiley: New York. p. 103-124.
15. Bradford, S.A. and R.W. Harvey, *Future research needs involving pathogens in groundwater*. Hydrogeology Journal, 2017. **25**(4): p. 931-938.
16. Harvey, R.W., *Microorganisms as tracers in groundwater injection and recovery experiments: a review*. Vol. 20. 1997. 461-472.
17. Sen, T.K., *Processes in Pathogenic Biocolloidal Contaminants Transport in Saturated and Unsaturated Porous Media: A Review*. Water, Air, & Soil Pollution, 2011. **216**(1-4): p. 239-256.
18. Taylor, R., A. Cronin, S. Pedley, J. Barker, and T. Atkinson, *The implications of groundwater velocity variations on microbial transport and wellhead protection – review of field evidence*. Vol. 49. 2004. 17-26.
19. Mac Kenzie, W.R., N.J. Hoxie, M.E. Proctor, M.S. Gradus, K.A. Blair, D.E. Peterson, J.J. Kazmierczak, D.G. Addiss, K.R. Fox, J.B. Rose, and J.P. Davis, *A Massive Outbreak in Milwaukee of Cryptosporidium Infection Transmitted through the Public Water Supply*. New England Journal of Medicine, 1994. **331**(3): p. 161-167.
20. O'Connor, D.R., *Part One Report of the Walkerton Inquiry: The Events of May 2000 and Related Issues*, O.M.o.t.A. General, Editor. 2002, Queen's Printer for Ontario, 2002.
21. Gallay, A., H. De Valk, M. Cournot, B. Ladeuil, C. Hemery, C. Castor, F. Bon, F. Mégraud, P. Le Cann, and J.C. Desenclos, *A large multi-pathogen waterborne community outbreak linked to faecal contamination of a groundwater system, France, 2000*. Clinical Microbiology and Infection, 2006. **12**(6): p. 561-570.
22. Fong, T.-T., L.S. Mansfield, D.L. Wilson, D.J. Schwab, S.L. Molloy, and J.B. Rose, *Massive Microbiological Groundwater Contamination Associated with a Waterborne Outbreak in Lake Erie, South Bass Island, Ohio*. Environmental Health Perspectives, 2007. **115**(6): p. 856-864.
23. Miettinen, I.T., O. Zacheus, C.H. von Bonsdorff, and T. Vartiainen, *Waterborne epidemics in Finland in 1998-1999*. Water Science and Technology, 2001. **43**(12): p. 67.
24. Wallender, E.K., E.C. Ailes, J.S. Yoder, V.A. Roberts, and J.M. Brunkard, *Contributing Factors to Disease Outbreaks Associated with Untreated Groundwater*. Groundwater, 2014. **52**(6): p. 886-897.
25. Reimus, P.W., M. Zavarin, and Y. Wang, *Colloid-Facilitated Radionuclide Transport: Current State of Knowledge from a Nuclear Waste Repository Risk Assessment Perspective*. 2017, Los Alamos National Laboratory (LANL).
26. McCarthy, J.F. and J.M. Zachara, *Subsurface transport of contaminants*. Environmental Science & Technology, 1989. **23**(5): p. 496-502.
27. Ryan, J.N. and M. Elimelech, *Colloid mobilization and transport in groundwater*. Colloids and Surfaces A: Physicochemical and Engineering Aspects, 1996. **107**(0): p. 1-56.
28. Ouyang, Y., D. Shinde, R.S. Mansell, and W. Harris, *Colloid-enhanced transport of chemicals in subsurface environments: A review*. Critical Reviews in Environmental Science and Technology, 1996. **26**(2): p. 189-204.
29. de Jonge, H., O.H. Jacobsen, L.W. de Jonge, and P. Moldrup, *Particle-Facilitated Transport of Prochloraz in Undisturbed Sandy Loam Soil Columns*. J. Environ. Qual., 1998. **27**(6): p. 1495-1503.
30. de Jonge, L.W., C. Kjaergaard, and P. Moldrup, *Colloids and Colloid-Facilitated Transport of Contaminants in Soils*. Vadose Zone J., 2004. **3**(2): p. 321-325.

31. Cho, K., X. Wang, S. Nie, Z. Chen, and D.M. Shin, *Therapeutic Nanoparticles for Drug Delivery in Cancer*. Clinical Cancer Research, 2008. **14**(5): p. 1310-1316.
32. Brigger, I., C. Dubernet, and P. Couvreur, *Nanoparticles in cancer therapy and diagnosis*. Advanced Drug Delivery Reviews, 2012. **64**: p. 24-36.
33. Hughes, D., *Transvascular fluid dynamics*. Veterinary Anaesthesia and Analgesia, 2000. **27**(1): p. 63-69.
34. Bradford, S.A., V.L. Morales, W. Zhang, R.W. Harvey, A.I. Packman, A. Mohanram, and C. Welty, *Transport and Fate of Microbial Pathogens in Agricultural Settings*. Critical Reviews in Environmental Science and Technology, 2012. **43**(8): p. 775-893.
35. Petosa, A.R., D.P. Jaisi, I.R. Quevedo, M. Elimelech, and N. Tufenkji, *Aggregation and Deposition of Engineered Nanomaterials in Aquatic Environments: Role of Physicochemical Interactions*. Environmental Science & Technology, 2010. **44**(17): p. 6532-6549.
36. Tufenkji, N., *Modeling microbial transport in porous media: Traditional approaches and recent developments*. Advances in Water Resources, 2007. **30**(6-7): p. 1455-1469.
37. Bradford, S.A. and S. Torkzaban, *Colloid Transport and Retention in Unsaturated Porous Media: A Review of Interface-, Collector-, and Pore-Scale Processes and Models* All rights reserved. No part of this periodical may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, or any information storage and retrieval system, without permission in writing from the publisher. Vadose Zone Journal, 2008. **7**(2): p. 667-681.
38. Zhang, W., X. Tang, N. Weisbrod, and Z. Guan, *A review of colloid transport in fractured rocks*. Journal of Mountain Science, 2012. **9**(6): p. 770-787.
39. McCarthy, J.F. and L.D. McKay, *Colloid Transport in the Subsurface*. Vadose Zone Journal, 2004. **3**(2): p. 326-337.
40. Molnar, I.L., W.P. Johnson, J.I. Gerhard, C.S. Willson, and D.M. O'Carroll, *Predicting colloid transport through saturated porous media: A critical review* Water Resources Research, 2015. **51**(50th anniversary special collection): p. 6804-6845.
41. Olivier, J., J. Vaxelaire, and E. Vorobiey, *Modelling of Cake Filtration: An Overview*. Separation Science and Technology, 2007. **42**(8): p. 1667-1700.
42. Lee, D.J. and C.H. Wang, *Theories of cake filtration and consolidation and implications to sludge dewatering*. Water Research, 2000. **34**(1): p. 1-20.
43. Tarleton, S. and R. Wakeman, *Solid/liquid separation: principles of industrial filtration*. 2005: Elsevier.
44. O'Carroll, D., B. Sleep, M. Krol, H. Boparai, and C. Kocur, *Nanoscale zero valent iron and bimetallic particles for contaminated site remediation*. Advances in Water Resources, 2013. **51**(0): p. 104-122.
45. Lin, D., X. Tian, F. Wu, and B. Xing, *Fate and Transport of Engineered Nanomaterials in the Environment* Journal of Environmental Quality, 2010. **39**(6): p. 1896-1908.
46. Pankhurst, Q.A., J. Connolly, S.K. Jones, and J. Dobson, *Applications of magnetic nanoparticles in biomedicine*. Journal of Physics D: Applied Physics, 2003. **36**(13): p. R167.
47. Wang, C.-S., *Electrostatic forces in fibrous filters—a review*. Powder Technology, 2001. **118**(1): p. 166-170.
48. Anderson, J.L., *Colloid transport by interfacial forces*. Annual review of fluid mechanics, 1989. **21**(1): p. 61-99.
49. Hermansson, M., *The DLVO theory in microbial adhesion*. Colloids and Surfaces B: Biointerfaces, 1999. **14**(1): p. 105-119.
50. Grasso, D., K. Subramaniam, M. Butkus, K. Strevett, and J. Bergendahl, *A review of non-DLVO interactions in environmental colloidal systems*. Reviews in Environmental Science and Biotechnology, 2002. **1**(1): p. 17-38.
51. Philippe, A. and G.E. Schaumann, *Interactions of Dissolved Organic Matter with Natural and Engineered Inorganic Colloids: A Review*. Environmental Science & Technology, 2014. **48**(16): p. 8946-8962.
52. Deichmann, U., *"Molecular" versus "colloidal": Controversies in biology and biochemistry, 1900–1940*. 2007.
53. Israelachvili, J.N., *11 - Contrasts between Intermolecular, Interparticle, and Intersurface Forces*, in *Intermolecular and Surface Forces (Third Edition)*, J.N. Israelachvili, Editor. 2011, Academic Press: San Diego. p. 205-222.
54. Pecora, R., *Dynamic Light Scattering Measurement of Nanometer Particles in Liquids*. Journal of Nanoparticle Research, 2000. **2**(2): p. 123-131.
55. Coffey, W.T. and Y.P. Kalmykov, *The Langevin equation: with applications to stochastic problems in physics, chemistry and electrical engineering*. 2004: World Scientific.
56. Nelson, K.E. and T.R. Ginn, *New collector efficiency equation for colloid filtration in both natural and engineered flow conditions*. Water Resources Research, 2011. **47**: p. 17.
57. Li, X.-q., D.W. Elliott, and W.-x. Zhang, *Zero-Valent Iron Nanoparticles for Abatement of Environmental Pollutants: Materials and Engineering Aspects*. Critical Reviews in Solid State and Materials Sciences, 2006. **31**(4): p. 111-122.
58. Wang, C.-B. and W.-x. Zhang, *Synthesizing Nanoscale Iron Particles for Rapid and Complete Dechlorination of TCE and PCBs*. Environmental Science & Technology, 1997. **31**(7): p. 2154-2156.
59. Carraway, E.R., A.J. Hoffman, and M.R. Hoffman, *Photocatalytic Oxidation of Organic Acids on Quantum-Sized Semiconductor Colloids*. Environmental Science & Technology, 1994. **28**(5): p. 786-793.
60. Klaine, S.J., P.J.J. Alvarez, G.E. Batley, T.F. Fernandes, R.D. Handy, D.Y. Lyon, S. Mahendra, M.J. McLaughlin, and J.R. Lead, *Nanomaterials in the environment: Behavior, fate, bioavailability, and effects*. Environmental Toxicology and Chemistry, 2008. **27**(9): p. 1825-1851.

- 695 61. Boparai, H.K., M. Joseph, and D.M. O'Carroll, *Kinetics and thermodynamics of cadmium ion removal by adsorption*
696 *onto nano zerovalent iron particles*. Journal of Hazardous Materials, 2011. **186**(1): p. 458-465.
- 697 62. Nichols, G., S. Byard, M.J. Bloxham, J. Botterill, N.J. Dawson, A. Dennis, V. Diart, N.C. North, and J.D. Sherwood, *A*
698 *review of the terms agglomerate and aggregate with a recommendation for nomenclature used in powder and particle*
699 *characterization*. Journal of Pharmaceutical Sciences, 2002. **91**(10): p. 2103-2109.
- 700 63. Bradford, S.A., J. Simunek, M. Bettahar, M.T. van Genuchten, and S.R. Yates, *Modeling Colloid Attachment,*
701 *Straining, and Exclusion in Saturated Porous Media*. Environmental Science & Technology, 2003. **37**(10): p. 2242-
702 2250.
- 703 64. Berkowitz, B., A. Cortis, M. Dentz, and H. Scher, *Modeling non-Fickian transport in geological formations as a*
704 *continuous time random walk*. Reviews of Geophysics, 2006. **44**(2): p. 49.
- 705 65. Netti, P.A., L.M. Hamberg, J.W. Babich, D. Kierstead, W. Graham, G.J. Hunter, G.L. Wolf, A. Fischman, Y. Boucher,
706 and R.K. Jain, *Enhancement of fluid filtration across tumor vessels: Implication for delivery of macromolecules*.
707 Proceedings of the National Academy of Sciences, 1999. **96**(6): p. 3137-3142.
- 708 66. Wadhams, G.H. and J.P. Armitage, *Making sense of it all: bacterial chemotaxis*. Nature Reviews Molecular Cell
709 Biology, 2004. **5**: p. 1024.
- 710 67. Ebel, J.P., J.L. Anderson, and D.C. Prieve, *Diffusiophoresis of latex particles in electrolyte gradients*. Langmuir, 1988.
711 **4**(2): p. 396-406.
- 712 68. Keh, H.J. and J.L. Anderson, *Boundary effects on electrophoretic motion of colloidal spheres*. Journal of Fluid
713 Mechanics, 2006. **153**: p. 417-439.
- 714 69. Yeap, S.P., A.L. Ahmad, B.S. Ooi, and J. Lim, *Electrosteric Stabilization and Its Role in Cooperative Magnetophoresis*
715 *of Colloidal Magnetic Nanoparticles*. Langmuir, 2012. **28**(42): p. 14878-14891.
- 716 70. Bakanov, S.P., *Thermophoresis of aerosols: a review*. Journal of Aerosol Science, 1991. **22**: p. S215-S218.
- 717 71. Paxton, W.F., S. Sundararajan, T.E. Mallouk, and A. Sen, *Chemical Locomotion*. Angewandte Chemie International
718 Edition, 2006. **45**(33): p. 5420-5429.
- 719 72. Buttinoni, I., J. Bialké, F. Kümmel, H. Löwen, C. Bechinger, and T. Speck, *Dynamical Clustering and Phase*
720 *Separation in Suspensions of Self-Propelled Colloidal Particles*. Physical Review Letters, 2013. **110**(23): p. 238301.
- 721 73. Korber, D.R., J.R. Lawrence, B. Sutton, and D.E. Caldwell, *Effect of laminar flow velocity on the kinetics of surface*
722 *recolonization by Mot+ and Mot- Pseudomonas fluorescens*. Microbial Ecology, 1989. **18**(1): p. 1-19.
- 723 74. Korber, D.R., J.R. Lawrence, and D.E. Caldwell, *Effect of Motility on Surface Colonization and Reproductive Success*
724 *of Pseudomonas fluorescens in Dual-Dilution Continuous Culture and Batch Culture Systems*.
725 Applied and Environmental Microbiology, 1994. **60**(5): p. 1421.
- 726 75. Yao, K.-M., M.T. Habibian, and C.R. O'Melia, *Water and waste water filtration. Concepts and applications*.
727 Environmental Science & Technology, 1971. **5**(11): p. 1105-1112.
- 728 76. Payatakes, A.C. and L. Gradoñ, *Dendritic deposition of aerosol particles in fibrous media by inertial impaction and*
729 *interception*. Chemical Engineering Science, 1980. **35**(5): p. 1083-1096.
- 730 77. Burganos, V.N., C.A. Paraskeva, and A.C. Payatakes, *Three-dimensional trajectory analysis and network simulation of*
731 *deep bed filtration*. Journal of Colloid and Interface Science, 1992. **148**(1): p. 167-181.
- 732 78. Payatakes, A.C., C. Tien, and R.M. Turian, *Part II. Case study of the effect of the dimensionless groups and*
733 *comparison with experimental data*. AIChE Journal, 1974. **20**(5): p. 900-905.
- 734 79. Payatakes, A.C., C. Tien, and R.M. Turian, *Trajectory calculation of particle deposition in deep bed filtration: Part I.*
735 *Model formulation*. AIChE Journal, 1974. **20**(5): p. 889-900.
- 736 80. Rajagopalan, R. and C. Tien, *TRAJECTORY ANALYSIS OF DEEP-BED FILTRATION WITH SPHERE-IN-CELL*
737 *POROUS-MEDIA MODEL*. Aiche Journal, 1976. **22**(3): p. 523-533.
- 738 81. Bhutra, S. and A.C. Payatakes, *Experimental investigation of dendritic deposition of aerosol particles*. Journal of
739 Aerosol Science, 1979. **10**(5): p. 445-464.
- 740 82. Payatakes, A.C. and L. Gradoñ, *Dendritic deposition of aerosols by convective Brownian diffusion for small,*
741 *intermediate and high particle Knudsen numbers*. AIChE Journal, 1980. **26**(3): p. 443-454.
- 742 83. Thomas, D., P. Penicot, P. Contal, D. Leclerc, and J. Vendel, *Clogging of fibrous filters by solid aerosol particles*
743 *Experimental and modelling study*. Chemical Engineering Science, 2001. **56**(11): p. 3549-3561.
- 744 84. Jiang, X., M. Tong, R. Lu, and H. Kim, *Transport and deposition of ZnO nanoparticles in saturated porous media*.
745 Colloids and Surfaces A: Physicochemical and Engineering Aspects, 2012. **401**(0): p. 29-37.
- 746 85. Wilhelm, C., F. Gazeau, J. Roger, J.N. Pons, and J.C. Bacri, *Interaction of Anionic Superparamagnetic Nanoparticles*
747 *with Cells: Kinetic Analyses of Membrane Adsorption and Subsequent Internalization*. Langmuir, 2002. **18**(21): p.
748 8148-8155.
- 749 86. Goodman, T.T., J. Chen, K. Matveev, and S.H. Pun, *Spatio-temporal modeling of nanoparticle delivery to multicellular*
750 *tumor spheroids*. Biotechnology and Bioengineering, 2008. **101**(2): p. 388-399.
- 751 87. Fujimori, K., D.G. Covell, J.E. Fletcher, and J.N. Weinstein, *A modeling analysis of monoclonal antibody percolation*
752 *through tumors: a binding-site barrier*. Journal of nuclear medicine : official publication, Society of Nuclear Medicine,
753 1990. **31**(7): p. 1191-1198.
- 754 88. Johnson, P.R. and M. Elimelech, *DYNAMICS OF COLLOID DEPOSITION IN POROUS-MEDIA - BLOCKING*
755 *BASED ON RANDOM SEQUENTIAL ADSORPTION*. Langmuir, 1995. **11**(3): p. 801-812.

89. Ma, H. and W.P. Johnson, *Colloid Retention in Porous Media of Various Porosities: Predictions by the Hemispheres-in-Cell Model*. Langmuir, 2010. **26**(3): p. 1680-1687.
90. Tufenkji, N. and M. Elimelech, *Correlation equation for predicting single-collector efficiency in physicochemical filtration in saturated porous media*. Environmental Science & Technology, 2004. **38**(2): p. 529-536.
91. Long, W. and M. Hilpert, *A Correlation for the Collector Efficiency of Brownian Particles in Clean-Bed Filtration in Sphere Packings by a Lattice-Boltzmann Method*. Environmental Science & Technology, 2009. **43**(12): p. 4419-4424.
92. Messina, F., D.L. Marchisio, and R. Sethi, *An extended and total flux normalized correlation equation for predicting single-collector efficiency*. Journal of Colloid and Interface Science, 2015. **446**: p. 185-193.
93. Kamai, T., M.K. Nassar, K.E. Nelson, and T.R. Ginn, *Colloid filtration prediction by mapping the correlation-equation parameters from transport experiments in porous media*. Water Resources Research, 2015. **51**(11): p. 8995-9012.
94. Lee, K.W. and B.Y.H. Liu, *Theoretical Study of Aerosol Filtration by Fibrous Filters*. Aerosol Science and Technology, 1982. **1**(2): p. 147-161.
95. Paraskeva, C.A., V.N. Burganos, and A.C. Payatakes, *THREE-DIMENSIONAL TRAJECTORY ANALYSIS OF PARTICLE DEPOSITION IN CONSTRICTED TUBES*. Chemical Engineering Communications, 1991. **108**(1): p. 23-48.
96. Johnson, W.P. and M. Hilpert, *Upscaling colloid transport and retention under unfavorable conditions: Linking mass transfer to pore and grain topology*. Water Resources Research, 2013. **49**(9): p. 5328-5341.
97. Logan, B., D. Jewett, R. Arnold, E. Bouwer, and C. O'Melia, *Clarification of Clean-Bed Filtration Models*. Journal of Environmental Engineering, 1995. **121**(12): p. 869-873.
98. Boccardo, G., E. Crevacore, R. Sethi, and M. Icardi, *A robust upscaling of the effective particle deposition rate in porous media*. Journal of Contaminant Hydrology, 2018. **212**: p. 3-13.
99. Nelson, K.E. and T.R. Ginn, *Colloid Filtration Theory and the Happel Sphere-in-Cell Model Revisited with Direct Numerical Simulation of Colloids*. Langmuir, 2005. **21**(6): p. 2173-2184.
100. Ma, H., E. Pazmino, and W.P. Johnson, *Surface Heterogeneity on Hemispheres-in-Cell Model Yields All Experimentally-Observed Non-Straining Colloid Retention Mechanisms in Porous Media in the Presence of Energy Barriers*. Langmuir, 2011. **27**(24): p. 14982-14994.
101. Ma, H., J. Pedel, P. Fife, and W.P. Johnson, *Hemispheres-in-Cell Geometry to Predict Colloid Deposition in Porous Media*. Environmental Science & Technology, 2009. **43**(22): p. 8573-8579.
102. Long, W., H. Huang, J. Serlemitsos, E. Liu, A.H. Reed, and M. Hilpert, *Pore-scale study of the collector efficiency of nanoparticles in packings of nonspherical collectors*. Colloids and Surfaces A: Physicochemical and Engineering Aspects, 2010. **358**(1-3): p. 163-171.
103. Elimelech, M. and C.R. Omelia, *KINETICS OF DEPOSITION OF COLLOIDAL PARTICLES IN POROUS-MEDIA*. Environmental Science & Technology, 1990. **24**(10): p. 1528-1536.
104. Elimelech, M. and C.R. Omelia, *EFFECT OF PARTICLE-SIZE ON COLLISION EFFICIENCY IN THE DEPOSITION OF BROWNIAN PARTICLES WITH ELECTROSTATIC ENERGY BARRIERS*. Langmuir, 1990. **6**(6): p. 1153-1163.
105. Hahn, M.W. and C.R. O'Melia, *Deposition and reentrainment of Brownian particles in porous media under unfavorable chemical conditions: Some concepts and applications*. Environmental Science & Technology, 2004. **38**(1): p. 210-220.
106. Elimelech, M., *Predicting collision efficiencies of colloidal particles in porous media*. Water Research, 1992. **26**(1): p. 1-8.
107. Shen, C., Y. Huang, B. Li, and Y. Jin, *Predicting attachment efficiency of colloid deposition under unfavorable attachment conditions*. Water Resources Research, 2010. **46**(11): p. W11526.
108. Bai, R. and C. Tien, *A New Correlation for the Initial Filter Coefficient under Unfavorable Surface Interactions*. Journal of Colloid and Interface Science, 1996. **179**(2): p. 631-634.
109. Bai, R. and C. Tien, *Particle Deposition under Unfavorable Surface Interactions*. Journal of Colloid and Interface Science, 1999. **218**(2): p. 488-499.
110. Vaidyanathan, R. and C.H.I. Tien, *HYDROSOL DEPOSITION IN GRANULAR BEDS—AN EXPERIMENTAL STUDY*. Chemical Engineering Communications, 1989. **81**(1): p. 123-144.
111. Chang, Y.-I.M. and J.-J. Whang, *Deposition of Brownian particles in the presence of energy barriers of DLVO theory: effect of the dimensionless groups*. Chemical Engineering Science, 1998. **53**(23): p. 3923-3939.
112. Kim, C. and S. Lee, *Effect of seepage velocity on the attachment efficiency of TiO₂ nanoparticles in porous media*. Journal of Hazardous Materials, 2014. **279**: p. 163-168.
113. Goldberg, E., M. Scheringer, T.D. Bucheli, and K. Hungerbühler, *Critical Assessment of Models for Transport of Engineered Nanoparticles in Saturated Porous Media*. Environmental Science & Technology, 2014. **48**(21): p. 12732-12741.
114. Wang, D., L. Ge, J. He, W. Zhang, D.P. Jaisi, and D. Zhou, *Hyperexponential and nonmonotonic retention of polyvinylpyrrolidone-coated silver nanoparticles in an Ultisol*. Journal of Contaminant Hydrology, 2014. **164**(0): p. 35-48.
115. Li, X.Q. and W.P. Johnson, *Nonmonotonic Variations in Deposition Rate Coefficients of Microspheres in Porous Media under Unfavorable Deposition Conditions*. Environmental Science & Technology, 2005. **39**(6): p. 1658-1665.
116. Tufenkji, N. and M. Elimelech, *Deviation from the classical colloid filtration theory in the presence of repulsive DLVO interactions*. Langmuir, 2004. **20**(25): p. 10818-10828.

117. Tufenkji, N., G.F. Miller, J.N. Ryan, R.W. Harvey, and M. Elimelech, *Transport of Cryptosporidium Oocysts in Porous Media: Role of Straining and Physicochemical Filtration*. Environmental Science & Technology, 2004. **38**(22): p. 5932-5938.
118. Tufenkji, N. and M. Elimelech, *Spatial Distributions of Cryptosporidium Oocysts in Porous Media: Evidence for Dual Mode Deposition*. Environmental Science & Technology, 2005. **39**(10): p. 3620-3629.
119. Kasel, D., S.A. Bradford, J. Šimůnek, M. Heggen, H. Vereecken, and E. Klumpp, *Transport and retention of multi-walled carbon nanotubes in saturated porous media: Effects of input concentration and grain size*. Water Research, 2013. **47**(2): p. 933-944.
120. Lanphere, J.D., C.J. Luth, and S.L. Walker, *Effects of Solution Chemistry on the Transport of Graphene Oxide in Saturated Porous Media*. Environmental Science & Technology, 2013. **47**(9): p. 4255-4261.
121. Li, Y., Y. Wang, K.D. Pennell, and L.M. Abriola, *Investigation of the Transport and Deposition of Fullerene (C60) Nanoparticles in Quartz Sands under Varying Flow Conditions*. Environmental Science & Technology, 2008. **42**(19): p. 7174-7180.
122. Mattison, N.T., D.M. O'Carroll, R. Kerry Rowe, and E.J. Petersen, *Impact of Porous Media Grain Size on the Transport of Multi-walled Carbon Nanotubes*. Environmental Science & Technology, 2011. **45**(22): p. 9765-9775.
123. Liu, X.Y., D.M. O'Carroll, E.J. Petersen, Q.G. Huang, and C.L. Anderson, *Mobility of Multiwalled Carbon Nanotubes in Porous Media*. Environmental Science & Technology, 2009. **43**(21): p. 8153-8158.
124. Camesano, T.A., K.M. Unice, and B.E. Logan, *Blocking and ripening of colloids in porous media and their implications for bacterial transport*. Colloids and Surfaces A: Physicochemical and Engineering Aspects, 1999. **160**(3): p. 291-307.
125. Tong, M., H. Ma, and W.P. Johnson, *Funneling of Flow into Grain-to-grain Contacts Drives Colloid-Colloid Aggregation in the Presence of an Energy Barrier*. Environmental Science & Technology, 2008. **42**(8): p. 2826-2832.
126. Molnar, I.L., J.I. Gerhard, C.S. Willson, and D.M. O'Carroll, *The impact of immobile zones on the transport and retention of nanoparticles in porous media*. Water Resources Research, 2015.
127. Fontes, D.E., A.L. Mills, G.M. Hornberger, and J.S. Herman, *Physical and chemical factors influencing transport of microorganisms through porous media*. Applied and Environmental Microbiology, 1991. **57**(9): p. 2473-2481.
128. Hornberger, G.M., A.L. Mills, and J.S. Herman, *Bacterial transport in porous media: Evaluation of a model using laboratory observations*. Water Resources Research, 1992. **28**(3): p. 915-923.
129. Lindqvist, R., J.S. Cho, and C.G. Enfield, *A kinetic model for cell density dependent bacterial transport in porous media*. Water Resources Research, 1994. **30**(12): p. 3291-3299.
130. Tong, M., X. Li, C.N. Brow, and W.P. Johnson, *Detachment-Influenced Transport of an Adhesion-Deficient Bacterial Strain within Water-Reactive Porous Media*. Environmental Science & Technology, 2005. **39**(8): p. 2500-2508.
131. Molnar, I.L., D.M. O'Carroll, and J.I. Gerhard, *Impact of surfactant-induced wettability alterations on DNAPL invasion in quartz and iron oxide-coated sand systems*. Journal of Contaminant Hydrology, 2011. **119**(1-4): p. 1-12.
132. Bradford, S.A., S. Torkzaban, and J. Šimunek, *Modeling colloid transport and retention in saturated porous media under unfavorable attachment conditions*. Water Resources Research, 2011. **47**.
133. Shipley, R.J. and S.J. Chapman, *Multiscale Modelling of Fluid and Drug Transport in Vascular Tumours*. Bulletin of Mathematical Biology, 2010. **72**(6): p. 1464-1491.
134. Torkzaban, S., S.S. Tazehkand, S.L. Walker, and S.A. Bradford, *Transport and fate of bacteria in porous media: Coupled effects of chemical conditions and pore space geometry*. Water Resources Research, 2008. **44**(4).
135. Pazmino, E.F., J. Trauscht, B. Dame, and W.P. Johnson, *Power Law Size-Distributed Heterogeneity Explains Colloid Retention on Soda Lime Glass in the Presence of Energy Barriers*. Langmuir, 2014. **30**(19): p. 5412-5421.
136. Trauscht, J., E. Pazmino, and W.P. Johnson, *Prediction of Nanoparticle and Colloid Attachment on Unfavorable Mineral Surfaces Using Representative Discrete Heterogeneity*. Langmuir, 2015. **31**(34): p. 9366-9378.
137. Wenning, L.A. and R.M. Murphy, *Coupled cellular trafficking and diffusional limitations in delivery of immunotoxins to multicell tumor spheroids*. Biotechnology and Bioengineering, 1999. **62**(5): p. 562-575.
138. Landkamer, L.L., R.W. Harvey, T.D. Scheibe, and J.N. Ryan, *Colloid transport in saturated porous media: Elimination of attachment efficiency in a new colloid transport model*. Water Resources Research, 2013. **49**(5): p. 2952-2965.
139. Schijven, J.F. and S.M. Hassanizadeh, *Virus removal by soil passage at field scale and groundwater protection of sandy aquifers (vol 46, pg 123, 2002)*. Water Science and Technology, 2002. **46**(6-7): p. 411-411.
140. Ginn, T.R., *On the distribution of multicomponent mixtures over generalized exposure time in subsurface flow and reactive transport: Theory and formulations for residence-time-dependent sorption/desorption with memory*. Water Resources Research, 2000. **36**(10): p. 2885-2893.
141. Contal, P., J. Simao, D. Thomas, T. Frising, S. Callé, J.C. Appert-Collin, and D. Bémer, *Clogging of fibre filters by submicron droplets. Phenomena and influence of operating conditions*. Journal of Aerosol Science, 2004. **35**(2): p. 263-278.
142. Frising, T., D. Thomas, D. Bémer, and P. Contal, *Clogging of fibrous filters by liquid aerosol particles: Experimental and phenomenological modelling study*. Chemical Engineering Science, 2005. **60**(10): p. 2751-2762.
143. Payet, S., D. Boulaud, G. Madelaine, and A. Renoux, *Penetration and pressure drop of a HEPA filter during loading with submicron liquid particles*. Journal of Aerosol Science, 1992. **23**(7): p. 723-735.

- 879 144. Agranovski, I.E. and R.D. Braddock, *Filtration of liquid aerosols on nonwetable fibrous filters*. AIChE Journal, 1998.
880 44(12): p. 2784-2790.
- 881 145. Cortis, A., T. Harter, L. Hou, E.R. Atwill, A.I. Packman, and P.G. Green, *Transport of Cryptosporidium parvum in*
882 *porous media: Long-term elution experiments and continuous time random walk filtration modeling*. Water Resources
883 Research, 2006. 42(12): p. n/a-n/a.
- 884 146. Elimelech, M., G. John, and J. Xiadong, *Particle deposition and aggregation: measurement, modelling and simulation*.
885 2013: Butterworth-Heinemann.
- 886 147. Kobayashi, M., H. Nanaumi, and Y. Muto, *Initial deposition rate of latex particles in the packed bed of zirconia beads*.
887 Colloids and Surfaces A: Physicochem. Eng. Aspects, 2009. 347: p. 2-7.
- 888 148. Ninham, B.W., *On progress in forces since the DLVO theory*. Advances in colloid and interface science, 1999. 83(1-3):
889 p. 1-17.
- 890 149. Van der Waals, J.D., *Over de Continuïteit van den Gas-en Vloeistoftoestand*. Vol. 1. 1873: Sijthoff.
- 891 150. Israelachvili, J., *Intermolecular and Surface Forces*. 1992, Academic Press: San Diego (CA).
- 892 151. Assemi, S., J. Nalaskowski, and P.W. Johnson, *Direct force measurements between carboxylate-modified latex*
893 *microspheres and glass using atomic force microscopy*. Colloids and Surfaces A: Physicochem. Eng. Aspects, 2006.
894 286: p. 70-77.
- 895 152. Foke, D.L., N.D. Prabhu, and J.A. Mann, *Formulation of the Short-Range Repulsion between Spherical Colloidal*
896 *Particle*. J. Phys. Chem. B, 1984. 88(23): p. 5735-5739.
- 897 153. Ohshima, H., *Electrokinetics of Soft Particles*. Colloid Polym. Sci., 2007. 285(13): p. 1411 - 1421.
- 898 154. Chandler, D., *Interfaces and the Driving Force of Hydrophilic Assembly*. Nature, 2005. 437: p. 640-647
- 899 155. Yang, X., S. Lin, and M.R. Wiesner, *Influence of natural organic matter on transport and retention of polymer coated*
900 *silver nanoparticles in porous media*. Journal of hazardous materials, 2014. 264: p. 161-168.
- 901 156. Dunphy Guzman, K.A., M.P. Finnegan, and J.F. Banfield, *Influence of Surface Potential on Aggregation and*
902 *Transport of Titania Nanoparticles*. Environmental Science & Technology, 2006. 40(24): p. 7688-7693.
- 903 157. Pensini, E., C. M. Yip, D. O'Carroll, and B.E. Sleep, *Carboxymethyl cellulose binding to mineral substrates:*
904 *Characterization by atomic force microscopy-based Force spectroscopy and quartz-crystal microbalance with*
905 *dissipation monitoring*. Journal of colloid and interface science, 2013. 402: p. 58-67.
- 906 158. Pensini, E., C. M. Yip, D.M. O'Carroll, and B.E. Sleep, *Effect of Water Chemistry and Aging on Iron-Mica Interaction*
907 *Forces: Implications for Iron Particle Transport*. Langmuir, 2012. 28(28): p. 10453-10463.
- 908 159. Pensini, E., C. M. Yip, D.M. O'Carroll, and B.E. Sleep, *Forces of interaction between fresh iron particles and iron*
909 *oxide (magnetite): Effect of water chemistry and polymer coatings*. Colloids and Surfaces A, 2013. 433 p. 104-110.
- 910 160. Pensini, E., C. M. Yip, D.M. O'Carroll, and B.E. Sleep, *Forces of interactions between iron and aluminum silicates:*
911 *Effect of water chemistry and polymer coatings*. Journal of colloid and interface science, 2013. 411: p. 8-15.
- 912 161. Pensini, E., B.E. Sleep, C.M. Yip, and D. O'Carroll, *Forces of interactions between bare and polymer-coated iron and*
913 *silica: Effect of ph, ionic strength, and humic acids*. Environmental science & technology, 2012. 46(24): p. 13401-
914 13408.
- 915 162. Sen, T.K. and K.C. Khilar, *Review on subsurface colloids and colloid-associated contaminant transport in saturated*
916 *porous media*. Advances in Colloid and Interface Science, 2006. 119(2-3): p. 71-96.
- 917 163. Torkzaban, S., S.A. Bradford, and S.L. Walker, *Resolving the coupled effects of hydrodynamics and DLVO forces on*
918 *colloid attachment in porous media*. Langmuir, 2007. 23(19): p. 9652-9660.
- 919 164. Redman, J.A., S.L. Walker, and M. Elimelech, *Bacterial adhesion and transport in porous media: Role of the*
920 *secondary energy minimum*. Environmental Science & Technology, 2004. 38(6): p. 1777-1785.
- 921 165. Israelachvili, J. and H. Wennerström, *Role of hydration and water structure in biological and colloidal interactions*.
922 Nature, 1996. 379(6562): p. 219.
- 923 166. Zhao, W., S.L. Walker, Q. Huang, and P. Cai, *Adhesion of bacterial pathogens to soil colloidal particles: influences of*
924 *cell type, natural organic matter, and solution chemistry*. Water research, 2014. 53: p. 35-46.
- 925 167. Hagens, W.I., A. G. Oomen, W.H. de Jong, F.R. Cassee, and A. Sips, *What do we (need to) know about the kinetic*
926 *properties of nanoparticles in the body?* Regulatory toxicology and pharmacology 2007. 49(3): p. 217-229.
- 927 168. Stylianopoulos, T., M.-Z. Poh, N. Insin, M.G. Bawendi, D. Fukumura, L.L. Munn, and R.K. Jain, *Diffusion of particles*
928 *in the extracellular matrix: the effect of repulsive electrostatic interactions*. Biophysical journal, 2010. 99(5): p. 1342-
929 1349.
- 930 169. Nel, A.E., L. Mädler, D. Velegol, T. Xia, E.M. Hoek, P. Somasundaran, F. Klaessig, V. Castranova, and M. Thompson,
931 *Understanding biophysicochemical interactions at the nano-bio interface*. Nature materials, 2009. 8(7): p. 543.
- 932 170. Phillips, R.J., W.M. Deen, and J.F. Brady, *Hindered transport in fibrous membranes and gels: effect of solute size and*
933 *fiber configuration*. Journal of colloid and interface science, 1990. 139(2): p. 363-373.
- 934 171. Jain, R.K., *Delivery of Novel Therapeutic Agents in Tumors: Physiological Barriers and Strategies*. JNCI: Journal of
935 the National Cancer Institute, 1989. 81(8): p. 570-576.
- 936 172. Alazraki, N.P., D. Eshima, L.A. Eshima, S.C. Herda, D.R. Murray, J.P. Vansant, and A.T. Taylor, *Lymphoscintigraphy,*
937 *the sentinel node concept, and the intraoperative gamma probe in melanoma, breast cancer, and other potential*
938 *cancers*. Seminars in Nuclear Medicine, 1997. 27(1): p. 55-67.

173. Krol, M.M., A.J. Oleniuk, C.M. Kocur, B.E. Sleep, P. Bennett, Z. Xiong, and D.M. O'Carroll, *A Field-Validated Model for In Situ Transport of Polymer-Stabilized nZVI and Implications for Subsurface Injection*. Environmental Science & Technology, 2013. **47**(13): p. 7332-7340.
174. Vogel, T.M., *Bioaugmentation as a soil bioremediation approach*. Current Opinion in Biotechnology, 1996. **7**(3): p. 311-316.
175. Bento, F.M., F.A.O. Camargo, B.C. Okeke, and W.T. Frankenberger, *Comparative bioremediation of soils contaminated with diesel oil by natural attenuation, biostimulation and bioaugmentation*. Bioresource Technology, 2005. **96**(9): p. 1049-1055.
176. Aderibigbe, A., K. Cheng, Z. Heidari, J. Killough, T. Fuss, and T. Stephens. *Detection of propping agents in fractures using magnetic susceptibility measurement enhanced by magnetic nano-particles*. in *SPE Annual Technical Conference and Exhibition*. Society of Petroleum Engineers.
177. Cocuzza, M., C. Opirri, V. Rocca, and F. Verga, *Current and future nanotech applications in the oil industry*. American Journal of Applied Sciences, 2012. **9**(6): p. 784-793.
178. Hwang, C.C., L. Wang, W. Lu, G. Ruan, G.C. Kini, C. Xiang, E.L.G. Samuel, W. Shi, A. Kan, M. Wong, M.B. Tomson, and J. Tour, *Highly stable carbon nanoparticles designed for downhole hydrocarbon detection*. Energy & Environmental Science, 2012. **5**: p. 8304-8304.
179. Liu, Q., M. Dong, S. Ma, and Y. Tu, *Surfactant enhanced alkaline flooding for Western Canadian heavy oil recovery*. Colloids and Surfaces A: Physicochemical and Engineering Aspects, 2007. **293**(1): p. 63-71.
180. Phenrat, T., N. Saleh, K. Sirk, R.D. Tilton, and G.V. Lowry, *Aggregation and sedimentation of aqueous nanoscale zerovalent iron dispersions*. Environmental Science & Technology, 2007. **41**(1): p. 284-290.
181. Albanese, A. and W.C.W. Chan, *Effect of Gold Nanoparticle Aggregation on Cell Uptake and Toxicity*. ACS Nano, 2011. **5**(7): p. 5478-5489.
182. Oberdörster, G., V. Stone, and K. Donaldson, *Toxicology of nanoparticles: A historical perspective*. Nanotoxicology, 2007. **1**(1): p. 2-25.
183. Phenrat, T., T.C. Long, G.V. Lowry, and B. Veronesi, *Partial Oxidation ("Aging") and Surface Modification Decrease the Toxicity of Nanosized Zerovalent Iron*. Environmental Science & Technology, 2009. **43**(1): p. 195-200.
184. Meng, H., M. Xue, T. Xia, Z. Ji, D.Y. Tarn, J.I. Zink, and A.E. Nel, *Use of Size and a Copolymer Design Feature To Improve the Biodistribution and the Enhanced Permeability and Retention Effect of Doxorubicin-Loaded Mesoporous Silica Nanoparticles in a Murine Xenograft Tumor Model*. ACS Nano, 2011. **5**(5): p. 4131-4144.
185. Hotze, E.M., T. Phenrat, and G.V. Lowry, *Nanoparticle Aggregation: Challenges to Understanding Transport and Reactivity in the Environment*. Journal of Environmental Quality, 2010. **39**(6): p. 1909-1924.
186. Adachi, Y., *Dynamic aspects of coagulation and flocculation*. Advances in Colloid and Interface Science, 1995. **56**: p. 1-31.
187. Cowell, C., R. Li-In-On, and B. Vincent, *Reversible flocculation of sterically-stabilised dispersions*. Journal of the Chemical Society, Faraday Transactions 1: Physical Chemistry in Condensed Phases, 1978. **74**: p. 337-347.
188. Perrault, S.D., C. Walkey, T. Jennings, H.C. Fischer, and W.C.W. Chan, *Mediating Tumor Targeting Efficiency of Nanoparticles Through Design*. Nano Letters, 2009. **9**(5): p. 1909-1915.
189. Sakulchaicharoen, N., D.M. O'Carroll, and J.E. Herrera, *Enhanced stability and dechlorination activity of pre-synthesis stabilized nanoscale FePd particles*. Journal of Contaminant Hydrology, 2010. **118**(3-4): p. 117-127.
190. Phenrat, T., A. Cihan, H.J. Kim, M. Mital, T. Illangasekare, and G.V. Lowry, *Transport and Deposition of Polymer-Modified Fe-0 Nanoparticles in 2-D Heterogeneous Porous Media: Effects of Particle Concentration, Fe-0 Content, and Coatings*. Environmental Science & Technology, 2010. **44**(23): p. 9086-9093.
191. Alili, L., M. Sack, A.S. Karakoti, S. Teuber, K. Puschmann, S.M. Hirst, C.M. Reilly, K. Zanger, W. Stahl, S. Das, S. Seal, and P. Brenneisen, *Combined cytotoxic and anti-invasive properties of redox-active nanoparticles in tumor-stroma interactions*. Biomaterials, 2011. **32**(11): p. 2918-2929.
192. Mornet, S., S. Vasseur, F. Grasset, and E. Duguet, *Magnetic nanoparticle design for medical diagnosis and therapy*. Journal of materials chemistry, 2004. **14**(14): p. 2161-2175.
193. Li, Q. and B.E. Logan, *Enhancing bacterial transport for bioaugmentation of aquifers using low ionic strength solutions and surfactants*. Water Research, 1999. **33**(4): p. 1090-1100.
194. Gentry, T., C. Rensing, and I.A.N. Pepper, *New Approaches for Bioaugmentation as a Remediation Technology*. Critical Reviews in Environmental Science and Technology, 2004. **34**(5): p. 447-494.
195. Hori, K. and S. Matsumoto, *Bacterial adhesion: From mechanism to control*. Biochemical Engineering Journal, 2010. **48**(3): p. 424-434.
196. Zhong, H., G. Liu, Y. Jiang, J. Yang, Y. Liu, X. Yang, Z. Liu, and G. Zeng, *Transport of bacteria in porous media and its enhancement by surfactants for bioaugmentation: A review*. Biotechnology Advances, 2017. **35**(4): p. 490-504.
197. Ryan, J.N., M. Elimelech, J.L. Baeseman, and R.D. Magelky, *Silica-Coated Titania and Zirconia Colloids for Subsurface Transport Field Experiments*. Environmental Science & Technology, 2000. **34**(10): p. 2000-2005.
198. Kocur, C.M., A.I. Chowdhury, N. Sakulchaicharoen, H.K. Boparai, K.P. Weber, P. Sharma, M.M. Krol, L. Austrins, C. Peace, B.E. Sleep, and D.M. O'Carroll, *Characterization of nZVI Mobility in a Field Scale Test*. Environmental Science & Technology, 2014. **48**(5): p. 2862-2869.

199. Velimirovic, M., T. Tosco, M. Uyttebroeck, M. Luna, F. Gastone, C. De Boer, N. Klaas, H. Sapon, H. Eisenmann, P.-O. Larsson, J. Braun, R. Sethi, and L. Bastiaens, *Field assessment of guar gum stabilized microscale zerovalent iron particles for in-situ remediation of 1,1,1-trichloroethane*. Journal of Contaminant Hydrology, 2014. **164**(0): p. 88-99.
200. Harvey, R.W., L.H. George, R.L. Smith, and D.R. LeBlanc, *Transport of microspheres and indigenous bacteria through a sandy aquifer: results of natural-and forced-gradient tracer experiments*. Environmental Science & Technology, 1989. **23**(1): p. 51-56.
201. Wilhelm, S., A.J. Tavares, Q. Dai, S. Ohta, J. Audet, H.F. Dvorak, and W.C.W. Chan, *Analysis of nanoparticle delivery to tumours*. Nature Reviews Materials, 2016. **1**: p. 16014.
202. Manzoor, A.A., L.H. Lindner, C.D. Landon, J.-Y. Park, A.J. Simnick, M.R. Dreher, S. Das, G. Hanna, W. Park, A. Chilkoti, G.A. Koning, T.L.M. ten Hagen, D. Needham, and M.W. Dewhirst, *Overcoming Limitations in Nanoparticle Drug Delivery: Triggered, Intravascular Release to Improve Drug Penetration into Tumors*. Cancer Research, 2012.
203. He, F., M. Zhang, T. Qian, and D. Zhao, *Transport of carboxymethyl cellulose stabilized iron nanoparticles in porous media: Column experiments and modeling*. Journal of Colloid and Interface Science, 2009. **334**(1): p. 96-102.
204. Tong, M. and W.P. Johnson, *Excess Colloid Retention in Porous Media as a Function of Colloid Size, Fluid Velocity, and Grain Angularity*. Environmental Science & Technology, 2006. **40**(24): p. 7725-7731.
205. Li, X.Q., P.F. Zhang, C.L. Lin, and W.P. Johnson, *Role of hydrodynamic drag on microsphere deposition and re-entrainment in porous media under unfavorable conditions*. Environmental Science & Technology, 2005. **39**(11): p. 4012-4020.
206. Torkzaban, S., S.A. Bradford, J.L. Vanderzalm, B.M. Patterson, B. Harris, and H. Prommer, *Colloid release and clogging in porous media: Effects of solution ionic strength and flow velocity*. Journal of Contaminant Hydrology, 2015. **181**: p. 161-171.
207. Quinn, J., C. Geiger, C. Clausen, K. Brooks, C. Coon, S. O'Hara, T. Krug, D. Major, W.-S. Yoon, A. Gavaskar, and T. Holdsworth, *Field Demonstration of DNAPL Dehalogenation Using Emulsified Zero-Valent Iron*. Environmental Science & Technology, 2005. **39**(5): p. 1309-1318.
208. Camesano, T.A. and B.E. Logan, *Influence of Fluid Velocity and Cell Concentration on the Transport of Motile and Nonmotile Bacteria in Porous Media*. Environmental Science & Technology, 1998. **32**(11): p. 1699-1708.
209. Bergendahl, J. and D. Grasso, *Prediction of colloid detachment in a model porous media: hydrodynamics*. Chemical Engineering Science, 2000. **55**(9): p. 1523-1532.
210. Flessner, M.F., J. Choi, K. Credit, R. Deverkadra, and K. Henderson, *Resistance of Tumor Interstitial Pressure to the Penetration of Intraperitoneally Delivered Antibodies into Metastatic Ovarian Tumors*. Clinical Cancer Research, 2005. **11**(8): p. 3117.
211. Chowdhury, A.I.A., M.M. Krol, C.M. Kocur, H.K. Boparai, K.P. Weber, B.E. Sleep, and D.M. O'Carroll, *nZVI injection into variably saturated soils: Field and modeling study*. Journal of Contaminant Hydrology, 2015. **183**: p. 16-28.
212. Henn, K.W. and D.W. Waddill, *Utilization of nanoscale zero-valent iron for source remediation—A case study*. Remediation Journal, 2006. **16**(2): p. 57-77.
213. Mondal, P.K., P.D. Furbacher, Z. Cui, M.M. Krol, and B.E. Sleep, *Transport of polymer stabilized nano-scale zero-valent iron in porous media*. Journal of Contaminant Hydrology, 2017.
214. DeFlaun, M.F., A.S. Tanzer, A.L. McAtter, B. Marshall, and S.B. Levy, *Development of an Adhesion Assay and Characterization of an Adhesion-Deficient Mutant of *Pseudomonas fluorescens**. Applied and Environmental Microbiology, 1990. **56**(1): p. 112-119.
215. DeFlaun, M.F., S.R. Oppenheimer, S. Streger, C.W. Condee, and M. Fletcher, *Alterations in Adhesion, Transport, and Membrane Characteristics in an Adhesion-Deficient Pseudomonad*. Applied and Environmental Microbiology, 1999. **65**(2): p. 759-765.
216. Streger, S.H., S. Vainberg, H. Dong, and P.B. Hatzinger, *Enhancing Transport of *Hydrogenophaga* and *flava* ENV735 for Bioaugmentation of Aquifers Contaminated with Methyl tert-Butyl Ether*. Applied and Environmental Microbiology, 2002. **68**(11): p. 5571.
217. Wang, W., W. Duan, S. Ahmed, A. Sen, and T.E. Mallouk, *From One to Many: Dynamic Assembly and Collective Behavior of Self-Propelled Colloidal Motors*. Accounts of Chemical Research, 2015. **48**(7): p. 1938-1946.
218. Paxton, W.F., A. Sen, and T.E. Mallouk, *Motility of Catalytic Nanoparticles through Self-Generated Forces*. Chemistry – A European Journal, 2005. **11**(22): p. 6462-6470.
219. Zheng, J.-m. and G.H. Pollack, *Long-range forces extending from polymer-gel surfaces*. Physical Review E, 2003. **68**(3): p. 031408.
220. Florea, D., S. Musa, J.M.R. Huyghe, and H.M. Wyss, *Long-range repulsion of colloids driven by ion exchange and diffusiophoresis*. Proceedings of the National Academy of Sciences of the United States of America, 2014. **111**(18): p. 6554-6559.
221. Machu, G., W. Meile, L.C. Nitsche, and U. Schaflinger, *Coalescence, torus formation and breakup of sedimenting drops: experiments and computer simulations*. J. Fluid Mech., 2001. **447**: p. 299-336.
222. Machu, G., L.N. W. M, and U. Schaflinger, *The motion of a swarm of particles traveling through a quiescent viscous fluid*. Zeitschrift Fur Angewandte Mathematik Und Mechanik, 2001. **81**: p. S547-S548.

- 1060 223. Adachi, K., S. Kiriya, and N. Koshioka, *The behaviour of a swarm of particles moving in a viscous fluid*. Chem.
1061 Eng. Sci., 1978. **33**: p. 115-121.
- 1062 224. Nitsche, J.M. and G.K. Batchelor, *Break-up of a falling drop containing dispersed particles*. Journal of Fluid
1063 Mechanics, 1997. **340**: p. 161-175.
- 1064 225. Schaflinger, U. and G. Machu, *Interfacial Phenomena in Suspensions*. Chemical Engineering Technology, 1999. **22**(7):
1065 p. 617-619.
- 1066 226. Metzger, B., M. Nicolas, and É. Guazzelli, *Falling clouds of particles in viscous fluids*. J. Fluid Mech., 2007. **580**: p.
1067 283-301.
- 1068 227. Pignatelli, F., M. Nicolas, and É. Guazzelli, *A falling cloud of particles at a small but finite Reynolds number*. J. Fluid
1069 Mech., 2011. **671**: p. 34-51.
- 1070 228. Lin, Y., J.H. Tan, N. Phan-Thien, and B.C. Khoo, *Settling of particle-suspension drops at low to moderate Reynolds
1071 numbers*. European Journal of Mechanics B-Fluids, 2017. **61**(1): p. 72-76.
- 1072 229. Rogers, W.B., *ART. XXXIII. --On the Formation of Rotating Rings by Air and Liquids*. American Journal of Science
1073 and Arts, 1858. **26**: p. 246-258.
- 1074 230. Tomlinson, C., *LXV. On a new variety of the cohesion-figures of liquids*. The London, Edinburgh, and Dublin
1075 Philosophical Magazine and Journal of Science, 1864. **27**(184): p. 425-432.
- 1076 231. Thomson, J.J. and H.F. Newall, *On the formation of vortex rings by drops falling into liquids, and some allied
1077 phenomena*. Proc. R. Soc. Lond., 1885. **39**: p. 417-435.
- 1078 232. Stucke, B., *Zur Bildung von Wirbelringen*. Zeitschrift für Physik, 1954. **137**: p. 376-379.
- 1079 233. Chapman, D.S. and P.R. Critchlow, *Formation of vortex rings from falling drops*. Journal of Fluid Mechanics, 1967.
1080 **29**(1): p. 177-185.
- 1081 234. Arecchi, F.T., P.K. Buah-Bassuah, F. Francini, C. Pérez-García, and F. Quercioli, *An experimental investigation of the
1082 break-up of a liquid drop falling in a miscible fluid*. Europhys. Lett., 1989. **9**: p. 333-338.
- 1083 235. Kojima, M., E.J. Hinch, and A. Acrivos, *The formation and expansion of a toroidal drop moving in a viscous fluid*.
1084 Phys. Fluids, 1984. **27**: p. 19-32.
- 1085 236. Subramanian, G. and D.L. Koch, *Evolution of clusters of sedimenting low-Reynolds-number particles with Oseen
1086 interactions*. Journal of Fluid Mechanics, 2008. **603**: p. 63-100.
- 1087 237. Happel, J. and H. Brenner, *Low Reynolds Number Hydrodynamics: With Special Applications to Particulate Media*.
1088 1983: Kluwer.
- 1089 238. Kim, S. and S.J. Karrila, *Microhydrodynamics: Principles and Selected Applications*. 2005: Dover.
- 1090 239. Barthès-Biesel, D., *Computer Simulation Using Particles*. 2012: CRC Press, New York.
- 1091 240. Boomsma, E.R., *Particle Swarms in Confining Geometries*. 2014, Purdue Univ. p. 104-104.
- 1092 241. Boomsma, E.R. and L.J. Pyrak-Nolte, *Particle Swarms in Fractures*. 2015, American Geophysical Union. p. 63-84.
- 1093 242. Bosse, T., L. Kleiser, J. Favre, and E. Meiburg, *Settling and breakup of suspension drops*. Physics of Fluids, 2005.
1094 **17**(9): p. 091107-091107.
- 1095 243. Northrup, E., *A photographic study of vortex rings in liquids*. Nature, 1912. **88**: p. 463-468.
- 1096 244. Mytyk, A., W. Meile, G. Brenn, and M.L. Ekiel-Jezewska, *Break-up of suspension drops settling under gravity in a
1097 viscous fluid close to a vertical wall*. Physics of Fluids, 2011. **23**(6): p. 1-15.
- 1098 245. Aranson, I.S., *Active colloids*. Phys. - Usp., 2013. **56**(1): p. 79-92.
- 1099 246. Di Leonardo, R., *Controlled collective motions*. Nat. Mater., 2016. **15**: p. 1057-1058.
- 1100 247. Ebbens, S.J., *Active colloids: Progress and challenges towards realising autonomous applications*. Curr. Opin. Colloid
1101 Interface Sci., 2016. **21**: p. 14-23.
- 1102 248. Zöttl, A. and H. Stark, *Emergent behavior in active colloids*. J. Phys.: Condens. Matter, 2016. **28**: p. 253001.
- 1103 249. Suarez, S.S. and A.A. Pacey, *Sperm transport in the female reproductive tract*. Hum Reprod Update, 2006. **12**(1): p.
1104 23-37.
- 1105 250. Marchiani, S., L. Tamburrino, M. Muratori, and E. Baldi, *Epididymal Sperm Transport and Fertilization*, in
1106 *Endocrinology of the Testis and Male Reproduction*, M. Simoni and I.T. Huhtaniemi, Editors. 2017, Springer
1107 International Publishing: Cham.
- 1108 251. Haig, S.J., G. Collins, R.L. Davies, C.C. Dorea, and C. Quince, *Biological aspects of slow sand filtration: past, present
1109 and future*. Water Sci. Tech. Water Supply, 2011. **11**(4): p. 468-472.
- 1110 252. Rippy and M. A., *Meeting the criteria: linking biofilter design to fecal indicator bacteria removal*. WIREs Water,
1111 2015. **2**: p. 577-592.
- 1112 253. Verma, S., A. Daverey, and A. Sharma, *Slow sand filtration for water and wastewater treatment – a review*.
1113 Environmental Technology Review, 2017. **6**(1): p. 47-58.
- 1114 254. Ginn, T.R., B.D. Wood, K.E. Nelson, T.D. Scheibe, E.M. Murphy, Clement, and T. Prabhakar, *Processes in microbial
1115 transport in the natural subsurface*. Adv. Water Resour., 2002. **25**(8-12): p. 1017-1042.
- 1116 255. Stevik, T.K., K. Aa, G. Ausland, and J.F. Hanssen, *Removal of Pathogenic Bacteria in Wastewater Percolating
1117 through Porous Media: A Review*. Water Research, 2004. **38**(6): p. 1355-1367.
- 1118 256. Kim, I.G., M.P. Hwang, P. Du, J. Ko, C.-w. Ha, S.H. Do, and K. Park, *Bioactive cell-derived matrices combined with
1119 polymer mesh scaffold for osteogenesis and bone healing*. Biomaterials, 2015. **50**: p. 75-86.
- 1120 257. Thursby, E. and N. Juge, *Introduction to the human gut microbiota*. Biochem J., 2017. **474** p. 1823-1836.

- 1121 258. Rolhion, N. and B. Chassaing, *When pathogenic bacteria meet the intestinal microbiota*. Phil. Trans. R. Soc. B, 2016.
1122 371: p. 20150504.
- 1123 259. Saintillan, D. and M.J. Shelley, *Orientational order and instabilities in suspensions of selflocomoting rods*. Phys. Rev.
1124 Lett., 2007. **99**: p. 058102.
- 1125 260. Baskaran, A. and M.C. Marchetti, *Statistical mechanics and hydrodynamics of bacterial suspensions*. Proc. Natl. Acad.
1126 Sci., 2009. **106**: p. 15567–15572.
- 1127 261. Hernandez-Ortiz, J.P., C.G. Stoltz, and M.D. Graham, *Transport and collective dynamics in suspensions of confined*
1128 *swimming particles*. Phys. Rev. Lett., 2005. **95**: p. 204501.
- 1129 262. Dreyfus, R., J. Baudry, M.L. Roper, M. Fermigier, H.A. Stone, and J. Bibette, *Microscopic artificial swimmers*. Nature,
1130 2005. **437**: p. 862.
- 1131 263. Wang, Y., R.M. Hernandez, D.J. Bartlett, J.M. Bingham, T.R. Kline, A. Sen, and T.E. Mallouk, *Bipolar*
1132 *electrochemical mechanism for the propulsion of catalytic nanomotors in hydrogen peroxide solutions*. Langmuir,
1133 2006. **22**: p. 10451–10456.
- 1134 264. Palacci, J., C. Cottin-Bizonne, C. Ybert, and B. L., *Sedimentation and effective temperature of active colloidal*
1135 *suspensions*. Phys. Rev. Lett., 2010. **105**: p. 088304.
- 1136 265. Wheat, P.M., N.A. Marine, J.L. Moran, and J.D. Posner, *Rapid fabrication of bimetallic spherical motors*. Langmuir,
1137 2010. **26**: p. 13052–13055.
- 1138 266. Tamm, S.L., *Ciliary Motion in Paramecium*. J Cell Biol, 1972. **55**: p. 250-255.
- 1139 267. Jana, S., A. Eddins, C. Spoon, and S. Jung, *Somersault of Paramecium in extremely confined environments*. Scientific
1140 Reports, 2015. **5**: p. 13148.
- 1141 268. Berg, H.C.E., *E. coli in motion*. 2004, New York: Springer-Verlag.
- 1142 269. Schwarz-Linek, J., J. Arlt, A. Jepsen, A. Dawson, T. Vissers, D. Miroli, T. Pilizota, V.A. Martinez, and W.C.K. Poon,
1143 *Escherichia coli as a model active colloid: A practical introduction*. Colloids Surf. B, 2016. **137**: p. 2-16.
- 1144 270. Frymier, P.D., R.M. Ford, H.C. Berg, and P.T. Cummings, *Three-dimensional tracking of motile bacteria near a solid*
1145 *planar surface*. Proc. Natl Acad. Sci. USA, 1995. **92**(13): p. 6195-6199.
- 1146 271. Lauga, E., W.R. DiLuzio, G.M. Whitesides, and H.A. Stone, *Swimming in circles: Motion of bacteria near solid*
1147 *boundaries*. . Biophysical Journal, 2006. **90**: p. 400–412.
- 1148 272. Berke, A.P., L. Turner, H.C. Berg, and E. Lauga, *Hydrodynamic attraction of swimming microorganisms by surfaces*.
1149 Phys. Rev. Lett., 2008. **101**: p. 038102.
- 1150 273. Li, G., J. Besson, L. Nisimova, D. Munger, P. Mahautmr, J.X. Tang, M.R. Maxey, and Y.V. Brun, *Accumulation of*
1151 *swimming bacteria near a solid surface*. Phys. Rev. E, 2011. **84**: p. 041932.
- 1152 274. Sartori, P., E. Chiarello, G. Jayaswal, M. Pierno, G. Mistura, P. Brun, A. Tiribocchi, and E. Orlandini, *Wall*
1153 *accumulation of bacteria with different motility patterns*. Phys. Rev. E, 2018. **97**(022610).
- 1154 275. Kantsler, V., J. Dunkel, M. Polin, and R. Goldstein, E., *Ciliary contact interactions dominate surface scattering of*
1155 *swimming eukaryotes*. Proc Natl Acad Sci USA, 2013. **110**(4): p. 1187–1192.
- 1156 276. Mirzakhani, M. and M.-R. Alam, *Flow characteristics of Chlamydomonas result in purely hydrodynamic scattering*.
1157 Phys. Rev. E, 2018. **98**: p. 012603.
- 1158 277. Takagi, D., J. Palacci, A.B. Braunschweig, M.J. Shelley, and J. Zhang, *Hydrodynamic capture of microswimmers into*
1159 *sphere-bound orbits*. Soft Matter, 2014. **10**: p. 1784-1789.
- 1160 278. Sipos, O., K. Nagy, R. Di Leonardo, and P. Galajda, *Hydrodynamic Trapping of Swimming Bacteria by Convex Walls*.
1161 Phys. Rev. Lett., 2015. **114**: p. 258104.
- 1162 279. Contino, M., E. Lushi, I. Tuval, V. Kantsler, and M. Polin, *Microalgae Scatter off Solid Surfaces by Hydrodynamic and*
1163 *Contact Forces*. Phys. Rev. Lett., 2015. **115**: p. 258102.
- 1164 280. Sklodowska, K., P.R. Debski, J.A. Michalski, P.M. Korczyk, M. Dolata, M. Zajac, and S. Jakiela, *Simultaneous*
1165 *Measurement of Viscosity and Optical Density of Bacterial Growth and Death in a Microdroplet*. Micromachines,
1166 2018. **9**: p. 251.
- 1167 281. Saintillan, D., *Rheology of Active Fluids*. Annual Review of Fluid Mechanics, 2018. **50**: p. 563–592.
- 1168 282. Zia, R.N., *Active and Passive Microrheology: Theory and Simulation*. Annual Review of Fluid Mechanics, 2018. **50**: p.
1169 371-405.
- 1170 283. Saintillan, D., *The dilute rheology of swimming suspensions: A simple kinetic model*. Exp. Mech., 2010. **50**: p. 1275.
- 1171 284. Ryan, S.D., B.M. Haines, L. Berlyand, F. Ziebert, and I.S. Aranson, *Viscosity of bacterial suspensions: Hydrodynamic*
1172 *interactions and self-induced noise*. . Phys. Rev. E., 2011. **83**: p. 050904.
- 1173 285. Rafai, S., L. Jibuti, and P. Peyla, *Effective viscosity of microswimmer suspensions*. Phys. Rev. Lett., 2010. **104**: p.
1174 098102.
- 1175 286. Mussler, M., S. Rafai, P. Peyla, and C. Wagner, *Effective viscosity of non-gravitactic Chlamydomonas Reinhardtii*
1176 *microswimmer suspensions*. EPL, 2013. **101**: p. 54004.
- 1177 287. López, H.M., J. Gachelin, C. Douarche, H. Auradou, and E. Clément, *Turning Bacteria Suspensions into Superfluids*.
1178 Phys. Rev. Lett., 2015. **115**: p. 028301.
- 1179 288. Takatori, S.C. and J.F. Brady, *Superfluid Behavior of Active Suspensions from Diffusive Stretching*. Phys. Rev. Lett.,
1180 2017. **118**: p. 018003.
- 1181 289. Clément, E., A. Lindner, C. Douarche, and H. Auradou, *Bacterial suspensions under flow*. Eur. Phys. J. Special Topics,
1182 2016: p. 2389-406.

- 1183 290. Hill, J., O. Kalkanci, J.L. McMurtry, and H. Koser, *Hydrodynamic surface interactions enable Escherichia coli to seek*
 1184 *efficient routes to swim upstream*. Phys. Rev. Lett., 2007. **98**: p. 068101.
- 1185 291. Figueroa-Morales, N., G.L. Miño, A. Rivera, R. Caballero, E. Clément, E. Altshuler, and A. Lindner, *Living on the*
 1186 *edge: transfer and traffic of E. coli in a confined flow*. Soft Matter, 2015. **11**: p. 6284-6293.
- 1187 292. Rusconi, R., J.S. Guasto, and R. Stocker, *Bacterial transport suppressed by fluid shear*. Nature Phys., 2014. **10**: p. 212-
 1188 217.
- 1189 293. Ezhilan, B. and D. Saintillan, *Transport of a dilute active suspension in pressure-driven channel flow*. J. Fluid Mech.,
 1190 2015. **777**: p. 482-522.
- 1191 294. Kim, M.K., F. Ingremeau, A. Zhao, B.L. Bassler, and H.A. Stone, *Local and global consequences of flow on bacterial*
 1192 *quorum sensing*. Nature Microbiology, 2018. **1**: p. 15005.
- 1193 295. Gaveau, A., C. Coetsier, C. Roques, P. Bacchin, and E. Dague, *Bacteria transfer by deformation through*
 1194 *microfiltration membrane*. Journal of Membrane Science, 2017. **523**: p. 446-455.
- 1195 296. Yazı, S.R., R. Nosrati, C.A. Stevens, D. Vogel, and C. Escobedo, *Migration of magnetotactic bacteria in porous media*.
 1196 Biomicrofluidics, 2018. **12**: p. 011101.
- 1197 297. Miño, G.L., M. Baabour, R. Chertcoff, G. Gutkind, E. Clément, H. Auradou, and I. Ippolito, *E coli Accumulation*
 1198 *behind an Obstacle*. Advances in Microbiology, 2018. **8**: p. 451-464.
- 1199 298. Nelson, K.E., A. Massoudieh, and T.R. Ginn, *E. Coli fate and transport in the Happel sphere-in-cell model*. Adv.
 1200 Water Resour., 2007. **30**(67): p. 1492:1504.

1201